
ADVANCES IN AQUACULTURE

2022

**Centre of Research for Advanced Aquaculture
(CORAA)**

*Edited by: Jiun-Yan Loh, Michelle Oi-Yoon Soo,
Baskaran Gunasekaran and Victor Charlie Andin*

 **UCSI University[®]**



Advances in Aquaculture

Edited by:

Jiun-Yan Loh, PhD

Michelle Oi-Yoon Soo, PhD

Baskaran Gunasekaran, PhD

Centre of Research for Advanced Aquaculture (CORAA),
UCSI University
Cheras, Kuala Lumpur, Malaysia.

and

Victor Charlie Andin, Msc

Marine Programme Conservation Department
World Wide Fund for Nature (WWF, Malaysia)
Petaling Jaya, Malaysia.

Preface

Global aquaculture has grown dramatically over the past 50 years and the trend is continually suppressing other animal production industry, making it the fastest-growing food production sector in the world. With huge seafood demands for a steadily growing population in 2050, the advancement of aquaculture production methods is vital. Modern farming practices and biotechnology are constantly being applied in the aquaculture sector to address various production limitations such as the supply of juveniles from hatcheries, nutritional quality, and etc.

This book “Advances in Aquaculture” provides a comprehensive, systematic guide to the use of current and emerging technologies in aquaculture production. The authors reviewed current technologies in algal production as a primary food source for fish and crustacean larvae. With the advancement of aquaculture technology, invertebrate production can be limitless in terms of stock quality and productivity. Nutrition is one of the major constraints in current aquaculture; some important nutritional issues emphasis in finfish and shellfish were discussed, the baseline information is of particular importance to the farmers. Last but not least, disease management is another biosecurity measure to ensure continuity and sustainability of aquaculture. With a range of upcoming and distinguished editors and expert contributors, “Advances in Aquaculture” is an authoritative review of the field for modern aquaculture.

This book would not have been possible without the dedicated efforts of the chapter authors. We hope this book will be able to stimulate research and better understanding of technological advances in growth, reproduction, and fish health management as more and more evidences show that aquaculture is the best animal production industry for protein and food to cater to the demand of a growing population in the world.

Jiun-Yan Loh
Centre of Research for Advanced Aquaculture
(CORAA)

Published in 2022 by UCSI Press

UCSI Heights, 1, Jalan Puncak Menara Gading,
Taman Connaught, 56000 Cheras,
Wilayah Persekutuan Kuala Lumpur

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted in any form, or by any means, electronic, mechanical, photocopying, recording or otherwise, without the prior permission of the publisher.

e ISBN 978-967-2782-28-5

Table of Contents

		Page
Chapter 1	Advances in Seaweed Cultivation with Focus on the Tropics	6
Chapter 2	Advances in Shrimp Aquaculture Technology	29
Chapter 3	Copepods as Live Feed for Marine Larviculture: Advantages, Challenges and Future Perspectives	38
Chapter 4	The Immunomodulatory Effects of Insect Meal in Aquaculture	47
Chapter 5	Latest Trends in Aquaculture Disease Management	61
Chapter 6	Current Update on the Strategies Used in Combating Acute Hepatopancreatic Necrosis Disease (AHPND)	77
Chapter 7	Insect-based Aquafeeds: A Sustainable Solution for Fish Nutrition	86

Chapter 1

ADVANCES IN SEAWEED CULTIVATION WITH FOCUS ON THE TROPICS

Hui-Yin Yeong^{1*}, *Swee-Sen Teo*^{2,3} & *Regina Zhi-Ling Leong*²

¹ Institute of Ocean and Earth Sciences (IOES), Universiti Malaya, 50603 Kuala Lumpur, Malaysia.

² Centre of Research for Advanced Aquaculture (CORAA), UCSI University, 56000, Cheras, Kuala Lumpur, Malaysia

³ Faculty of Applied Sciences, UCSI University, 56000 Kuala Lumpur, Malaysia.

*Corresponding author: yeong@um.edu.my

ABSTRACT

Over the last 50 years, the rapid development of seaweed cultivation technologies in tropical countries has resulted in the seaweed aquaculture industry producing more than 99% of seaweed biomass production in the tropics. In 2020, the major farmed seaweed species in the tropics, namely *Euचेuma* spp. (8,194,053.60 tonnes FW), *Gracilaria* spp. (5,176,744.42 tonnes FW), and *Kappaphycus alvarezii* (1,603,213.33 tonnes FW), accounted for 98.92%, 99.90%, and 98.92% of the respective species' global aquaculture production, with USD 5,653,007.86 in value. The main methods of growing seaweed were line cultivation, floating raft cultivation for *Euचेuma* spp. and *Kappaphycus* spp., and open water and pond cultivation for *Gracilaria* spp. These methods have been used in selected tropical countries since the 1970s and 1980s. However, there are still many improvements needed and countless challenges to overcome for the sustainable integration of seaweed cultivation production into the new era of the global aquaculture industry.

Keywords: Seaweed Cultivation, Algae Biotechnology, Seaweed Aquaculture, *Euचेuma*, *Gracilaria*, *Kappaphycus alvarezii*

1.1 Introduction

Seaweeds are benthic macroalgae present in marine habitats. Seaweeds are found along the shores of all areas, from the tropics to the polar regions, and play an important part in the ecology of the ocean. The macroalgae are thallophytes, which implies that they do not have well-defined vascular tissues, whilst lacking roots, stems, and leaf structures. Nevertheless, they do have chlorophyll as a major photosynthetic pigment. Seaweeds also do not possess complicated reproductive systems and do not have sterile cells in their reproductive systems, meaning that all cells are potentially fertile (Lee, 1999; Barsanti & Gualtieri, 2006, Baweja et. al. 2016). Seaweeds serve as primary producers in the food chain of the marine ecosystem. As such, they are an essential source of oxygen, carbon dioxide, and water for the metabolic processes of marine organisms. They produce organic materials from sunlight too (Lee, 1999; Baweja et. al. 2016). Seaweeds can be classified into three main groups, namely green seaweed (Chlorophyta), brown seaweed (Ochrophyta) and red seaweed (Rhodophyta) based on their characteristic pigmentation. There are approximately 1,800 species of green seaweed, 6,200 species of red seaweed, and 1,800 species of brown seaweed reported found in the natural environment (Mouritsen, 2013).

Today, seaweed is one of the marine resources that has received substantial interest, particularly in developing countries. This is due to the high demand for seaweed as a source of food for direct human consumption, animal feed in the agriculture and aquaculture industries, fertilizer and soil conditioner in the agriculture business, and a source of industrial components in the food and medical industries. Furthermore, because of the intensive development of algae biotechnology, seaweed has emerged as a source of bioactive natural compounds that may have value in the nutraceutical, cosmeceutical, and pharmaceutical industries. Seaweed is also a source of biomass that can be used in the production of alternative bioenergy and biofuels.

The increasing demand for seaweed resources as raw materials from various industries globally has led to an increase in world seaweed production and utilization. The annual global production of seaweed increased from 2,116,809 tonnes (fresh weight) in 1950 to 36,165,326 tonnes (fresh weight) in 2020 (FAO, 2022b), representing an increase of around 8879.87% over the course of the same period. This biomass of seaweed was collected either from natural populations or from the seaweed aquaculture industry. The rapidly increasing demand for seaweed biomass has also resulted in the expansion of the seaweed aquaculture business and the development of techniques for seaweed cultivation, both of which are intended to increase the amount of seaweed produced through seaweed aquaculture. This led to a rise in the number of interested parties and investors who invested in the development of advanced cultivation technologies for the seaweed aquaculture industry (García-Poza et. al. 2020).

1.2 Seaweed Production

The annual world seaweed production can be attributed to the harvesting of seaweed from both wild populations and farming industries. In 1950, the annual world seaweed production amounted to 407,273 tonnes fresh weight (FW), of which 91.50% (372,637 tonnes FW) was taken from the wild populations and 8.50% (34,636 tonnes FW) was harvested by the seaweed aquaculture industry. The annual production of seaweed had climbed to 2,116,809 tonnes FW by the year 1970. Of this total, 54.06% came from farmed seaweed (1,144,380 tonnes FW), while 45.94% was collected from the wild population (972,429 tonnes FW). Since 1970, there has been a substantial increase in the annual world seaweed production (Figure 1.1).

In 2020, world seaweed production achieved an annual output of 36,165,326 tonnes FW, of which 96.81% (35,013,088 tonnes FW) was produced by the seaweed aquaculture farming industry, and this aquaculture production was worth an annual value of USD 16,444,926,480. In 2020, the highest seaweed aquaculture production was *Laminaria japonica* (12,469,807 tonnes FW), and followed by *Euचेuma* spp (8,283,513 tonnes FW), *Gracilaria* spp (5,180,416 tonnes FW), *Undaria pinnatifida* (2,810,605 tonnes FW), *Porphyra* spp (2,220,180 tonnes FW) and *Kappaphycus alvarezii* (1,604,110 tonnes FW) (FAO, 2022b).




Figure 1.1 World seaweed production (1950 -2020) (Data extracted from FAO, 2022).

In the tropical region, the annual seaweed production started to increase greatly after the 1970s, when seaweed farming was introduced to the tropics. In 1950, the annual production of tropical seaweed was 17,060 tonnes FW, of which 94.31% (16,090 tonnes FW) was harvested from wild populations and 5.69% (970 tonnes FW) came from seaweed farms (Figure 1.2). In 1980, seaweed aquaculture production started to dominate the seaweed production in the tropics, contributing to 61.83% (143,150 tonnes FW) of the annual seaweed production in the tropics. Since 2005, tropical

seaweed aquaculture has contributed more than 95% of the annual seaweed production in the tropics. In 2020, 99.08 % (15,061,383 tonnes FW) of total annual seaweed production in the tropics was harvested by tropical seaweed aquaculture farming, with USD 5,681,861.48 in value (FAO, 2022a).




Figure 1.2 Tropical seaweed production (1950 -2020) (Data extracted from FAO, 2022).

In spite of the fact that tropical nations produced approximately 35.92% of worldwide seaweed aquaculture production in 2020, only three major seaweeds are actively cultivated in tropical regions. The annual tropical seaweed aquaculture production from 1985 to 2020 is summarized in Table 1.1. In 2020, the major farmed seaweed species in the tropics, namely *Eucheuma* spp. (8,194,053.60 tonnes FW), *Gracilaria* spp. (5,176,744.42 tonnes FW), and *Kappaphycus alvarezii* (1,603,213.33 tonnes FW), accounted for 98.92%, 99.90%, and 98.92% of the respective species' global aquaculture production. Table 1.2 summarizes the major seaweed production (tonnes (FW) and value (USD)) and the top three producing countries in the tropics.

Table 1.2 Major seaweed production (tonnes FW) and value (USD) in the Tropics (2000 and 2020)*.

No	Country*	2000		2020	
		Production (tonnes FW)	Value (USD)	Production (tonnes FW)	Value (USD)
1	<i>Eucheuma spp</i>				
	Indonesia	197,227.00	21,694.97	8,080,796.00	1,640,850.95
	Philippines	34,424.00	2,953.24	63,073.20	8,066.93
	China	0.00		38,560.00	32,351.84
	Total	249,730.00	25,449.03	8,194,053.60	1,684,999.44
2	<i>Gracilaria spp</i>				
	China	0.00	0.00	3,689,670.00	2,025,628.83
	Indonesia	8,000.00	880.00	1,456,730.00	117,279.94
	Chile	33,471.00	16,735.50	18,268.00	216,475.80
	Total	51,514.00	17,692.41	5,176,744.42	2,359,412.56
3	<i>Kappaphycus alvarezii</i>				
	Philippines	644,319.00	46,962.78	1,404,743.36	205,300.02
	Malaysia	0.00	0.00	182,061.00	1,400,419.70
	Solomon Islands	0.00	0.00	5,500.00	205.58
	Total	649,464.00	47,828.15	1,603,213.33	1,608,595.86
4	<i>Caulerpa spp</i>				
	Philippines	28,296.00	3,367.62	789.36	515.38
	Indonesia	0.00	0.00	232.00	107.96
	Total	28,296.00	3,367.62	1,021.36	623.34

* The top 3 seaweed producing countries in the tropics in 2020

**Data extracted from FAO. 2022. Fishery and Aquaculture Statistics. Global aquaculture production 1950-2020 (FishStatJ). In: FAO Fisheries and Aquaculture Division [online]. Rome. Updated 2022.

Table 1.1 Annual seaweed production (tonnes FW) and value (USD) in the Tropics (1985 - 2020)*.

No	Seaweed	1985	1990	1995	2000	2005	2010	2015	2020
1	<i>Caulerpa</i> spp (tonnes FW)	11,278.00	17,471.00	4,040.00	28,296.00	4,309.00	4,309.00	1,219.00	1,021.36
	<i>Caulerpa</i> spp (USD)	3,637.22	4,699.63	458.62	3,367.62	1,549.38	2,560.97	1,073.99	623.34
2	<i>Eucheuma</i> spp (tonnes FW)	9,668.00	26,939.00	19,245.00	249,730.00	1,089,116.00	3,611,634.40	10,295,622.60	8,194,053.60
	<i>Eucheuma</i> spp (USD)	1,211.49	3,183.11	3,263.51	25,449.03	166,896.05	1,147,173.44	806,508.91	1,684,999.44
3	<i>Gracilaria</i> spp (tonnes FW)	6,524.00	40,057.00	57,222.00	51,514.00	933,184.00	1,657,144.00	3,767,020.00	5,176,744.42
	<i>Gracilaria</i> spp (USD)	1,999.60	17,500.62	22,184.65	17,692.41	276,706.87	589,445.65	1,614,045.11	2,359,412.56
4	<i>Kappaphycus alvarezii</i> (tonnes FW)	162,385.00	259,247.00	562,043.00	649,464.00	1,279,234.80	1,884,024.10	1,746,008.30	1,603,213.33
	<i>Kappaphycus alvarezii</i> (USD)	28,236.23	37,373.63	53,452.63	47,828.15	107,167.66	266,368.59	212,654.87	1,608,595.86
5	Others (tonnes FW)	0.00	0.00	0.00	0.00	16,125.00	2,900.00	1,565.00	1,300.00
	Others (USD)	1,843.22	5,000.00	11,220.00	1,824.69	1,234.09	641.29	462.29	28,230.28
	Total (tonnes FW)	135,186.00	189,855.00	343,714.00	642,550.00	995,129.00	3,308,743.80	7,158,676.50	15,811,169.90
	Total (USD)	36,927.76	67,756.99	90,579.41	96,161.90	553,554.05	2,006,189.94	2,634,745.17	5,681,861.48

*Data extracted from FAO. 2022. Fishery and Aquaculture Statistics. Global aquaculture production 1950-2020 (FishStatJ). In: FAO Fisheries and Aquaculture Division [online]. Rome. Updated 2022.

1.3 Seaweed Utilization

Since 2500 BC, seaweed has been utilized in China (Tseng, 1981). However, Hoppe and Schmid (1969) state that the first few true instances of economic exploitation of seaweed were carried out in China and Japan around the year 1670. There have been at least 291 different types of seaweed utilized on a global scale up to this point. This includes 33 species of green seaweed, 75 species of brown seaweed, and 163 species of red seaweed, as reported by Lindsey White and Peter Wilson (2015) (Table 1.3). In 2019, the world seaweed aquaculture industry was cultivating a total of 27 different seaweed species (items as from ASFIS8 (database of the Aquatic Sciences and Fisheries Information System), but only 3 major seaweed groups or 5 seaweed species were cultivated in tropical regions (Cai et al, 2021), including *Euचेuma* spp., *Gracilaria* spp. and *Kappaphycus alvarezii*.

Table 1.3 Summary of the composition, important properties, uses, and values of the commercially important phycocolloids seaweeds (*Euचेuma* spp, *Gracilaria* spp., and *Kappaphycus alvarezii*)*.

Seaweeds	Uses and Applications
<p><i>Euचेuma</i> spp, <i>Kappaphycus</i> spp.</p>	<p>Carrageenans</p> <ul style="list-style-type: none"> • <i>Kappaphycus alvarezii</i> - important source of kappa carrageenan • <i>Euचेuma denticulatum</i> - important source of iota carrageenan <p>• Composition</p> <ul style="list-style-type: none"> ▪ Kappa- and iota- ▪ Alternating 1,3-linked α-D-galactosa and 1,4-linked 3,6-anhydro-β-D-galactosa backbone (carrabiose) substituted with varying percentages of ester sulphate. <p>• Important properties</p> <ul style="list-style-type: none"> ▪ Bind moisture. ▪ Stabilize emulsions ▪ Control flow and texture properties of food systems. ▪ High protein reactivity- strong interactions with milk proteins <p>• Selected application</p> <ul style="list-style-type: none"> ▪ dairy products: kappa acarrageenan added in cottage cheese/ ice cream to prevent separation & control texture ▪ Water-based foods: as substitutes for gelatin; suitable for vegetarians ▪ meat products: carrageenan binds water within the poultry muscle and improves texture and tenderness. ▪ Frozen dessert stabilizers. ▪ Chocolate milk stabilizer. ▪ Texturizers for low fat foods. ▪ Low-calories jellies. ▪ Toothpaste binders. ▪ Air-freshener gels. ▪ Personal care products. ▪ Immobilized biocatalysts.

Gracilaria spp.	<p>Agar</p> <ul style="list-style-type: none"> • Composition <ul style="list-style-type: none"> ▪ Alternating 1, 4-linked α-D galactose and 3,6 anhydro-α-1 galactose backbone (agarobiose) substituted with varying percentages of methoxyl, ester sulphate and ketal pyruvate groups • Important properties <ul style="list-style-type: none"> ▪ Gel aqueous at low concentrations. ▪ Form thermo reversible gels. ▪ Relatively inert. ▪ Significant degree to hysteresis. ▪ Retain moisture. ▪ Resist hydrolysis by terrestrial microorganisms. ▪ Belong to galatans. ▪ Give viscosity to aqueous solutions. • Selected application <ul style="list-style-type: none"> ▪ Dental impression media ▪ Jelly candies ▪ Baking icings ▪ Jams and soups ▪ Canned meat ▪ Laxatives ▪ Microbial culture media ▪ Raw material for agarose ▪ Textile sizing and block printing <p>Agarose</p> <ul style="list-style-type: none"> • Composition <ul style="list-style-type: none"> ▪ Alternating 1, 4-linked α-D galactose and 3,6 anhydro-α-1 galactose backbone (agarobiose) substituted with varying percentages of methoxyl, ester sulphate and ketal pyruvate groups • Important properties <ul style="list-style-type: none"> ▪ Gel aqueous at low concentrations. ▪ From ion-dependent thermo reversible gels. ▪ Controllable electroendosinosis. ▪ Minimal non-specific protein reactivity. ▪ Significant degree of hysteresis. • Selected application <ul style="list-style-type: none"> ▪ Material for: Electrophoresis, Immunoassays ▪ Microbial and cell culture ▪ Chromatography ▪ Immobilized system

*Modified from Jensen, 1993; Renn, 1997; McHugh, 2002; Phang et al 1996; Phang et al 2010a & 2010b; Bixler & Porse, 2010 and Phang, 2010)

Aside from that, seaweed has a greater potential for the production of bioplastics than bacterial and plant sources because it is extremely rich in polysaccharides such as carrageenan, agar, floridean, starch, and alginate. This is because seaweed is able to convert its starch into alginate. These chemicals provide adequate properties to make edible bioplastic films and packages with high tensile strength, flexibility, and elasticity. These qualities are desirable in bioplastics (Moey et al., 2015). The usage of kappa-carrageenan, which is derived from Rhodophyta seaweed, has been introduced as a precursor for the development of bioplastic film. This can improve the moisture barrier and tensile strength of conventional bioplastic films (Abdul Khalil et al., 2017). Conventional plastics that land up in landfills are the root cause of today's plastic epidemic and may be completely broken down into its component parts by bioplastics made from seaweed at a far faster rate. This may provide a sustainable solution to the problem that conventional plastic presents.

Due to the high concentration of rich polysaccharides and the gelling properties of seaweed, it is frequently used as the primary ingredient or as an additive in a variety of industries and applications, including the food industry, industrial applications, agriculture, skincare, and therapeutic fields (Ganesan et al., 2019; Cherry et al., 2019). Recent developments in water bioremediation have seen an increased emphasis placed on the utilization of natural materials or biomass. In order to find a suitable replacement for chemically-manufactured adsorbents, seaweed has been proposed as a possible biosorbent for the removal of pollutants in water bioremediation. These pollutants include textile dyes, heavy metals, and organic wastes (Oualid et al., 2020; D. Shah et al., 2021; Buhani et al., 2021). According to a study that was conducted by Abdullah Al-Dhabi and Arasu (2022), the phytoremediation process used by marine seaweed was able to effectively remove heavy metals from wastewater such as lead (Nessim et al., 2011; Eka Putri et al., 2018; Isam et al., 2019).

In recent research, seaweed has been utilized for surface modification in magnetic nanoparticle adsorbents in order to improve the adsorbent's stability. This was accomplished by increasing the surface area of the adsorbent. Research into the adsorption of proteins and the removal of dyes using magnetic nanoparticles with kappa-carrageenan as a surface modifier has been conducted (Mahdavinia & Etemadi, 2019; Salgueiro et al., 2013; Soares et al., 2017). The coating layer not only helps to improve the colloidal stability of the nanoparticles but also works as a binder to adsorb heavy metals. This is one of the many roles that the coating plays (Wan et al., 2021). Electrostatic attraction was one of the potential mechanisms that may be used to accomplish the goal of covering the positively charged surfaces of magnetic nanoparticles with negatively charged kappa-carrageenan. According to research, the nanoparticles that were synthesized improve the removal of heavy metals and organic contaminants from wastewater. This is accomplished by adsorbing the heavy metals to the surface-active sites of the adsorbent through the electrostatic interaction of the functional groups (Yew et al., 2016; Kulal & Badalamoole, 2020). Therefore, cultivation of seaweed has developed into a significant sector as a means of satisfying the high global demand for the seaweed industry and of compensating for the loss of seaweed populations because of climate change on a global scale.

1.4 Seaweed Cultivation Techniques

In general, the seaweed aquaculture sector in the tropics focuses on three of the most commercially important genera: *Eucheuma*, *Gracilaria*, and *Kappaphycus*. There are different approaches on the cultivation techniques for Eucheumatoids (which include those from the genera *Eucheuma* and *Kappaphycus*) (Figure 1.3) and *Gracilaria* spp.

In the early 1970s, eucheumatoid farming started in southern Philippines and gradually was introduced to the other tropical countries such as Indonesia, Malaysia, the Pacific Island, South Africa, Madagascar and Indo-China (Hurtado & Crithcley, 2006). Cultivation techniques include line cultivation (long line/ monoline cultivation) and floating raft cultivation techniques.



Figure 1.3 Some varieties of *Kappaphycus* culture are harvested from the seaweed farm in Sabah, Malaysia.

1.4.1 Line cultivation (long line/ monoline cultivation)

Basically, line cultivation (long line or monoline cultivation) is carried out in bays or open seas, away from sources of freshwater such as rivers or estuarine areas and other harmful sources such as industrial waste or agricultural waste. The site must also be shielded from large tidal and wind-generated waves to prevent destruction of the farm by these environmental disasters (Pereira & Yarish, 2008).

Selected eucheumatoid seedlings are tied to a short, thinner rope, which is then fastened to the main ropes at sea that range in length from 10 meters to 50 meters, or longer. These ropes are laid out in a parallel manner with variable distances between each other, about 0.5 m to 1.0 m or more, depending on the size of the seaweed that is going to be harvested. This rope, which is stuffed with seaweed, will be positioned at depths that range from those listed below:



Kappaphycus salad



Kerabu sare (*Gracilaria* salad)



Noodles made from seaweeds



Agar-agar powder for jelly



Personal care products made from seaweed



Seaweed soap



Seaweed snacks*



3-in-1 seaweed coffee

Figure 1.4 Some available products from *Eucheuma* spp, *Gracilaria* spp and *Kappaphycus*. *Photo credit: Liu Yung Kuan, Famous Alpine Seaweed Farm

- a) Off-bottom – The long planting rope will be positioned close to the bottom along the coast, ideally with at least around 0.3 meters of water on top of it at the lowest tide (Figure 1.5). This technique is typically utilized with species that are either small or frequently harvested, such as *Eucheuma* and *Kappaphycus*.



Figure 1.5 Off-bottom seaweeds cultivation model

- b) Submerged hanging line - The long planting rope will be positioned in the midwater near to the coast with the condition that it may be submerged in the water during high tides and that it may be at the surface of the water or even exposed during low tides.
- c) Long line/floating line - With the support of float balls or aquaculture buoys, the long planting rope will be positioned at or near the surface of the water; meanwhile, the ends of the two ends will be attached and maintained with anchors at the bottom. The cultivated seaweed that is tied along the long rope is kept submerged but is allowed to float relatively close to the seawater's surface (Figure 1.6 and 1.7). With this technique, it is possible to cultivate seaweed regardless of the depth of the seafloor.

1.4.2 Floating raft cultivation

This cultivation technique takes place on the near seawater surface, where the seaweed is attached to lines or nets within a floating rigid frame (Figure 1.8), made of bamboo or some other material.



Figure 1.6 Long line cultivation system at Semporna, Sabah, Malaysia.



Figure 1.7 *Kappaphycus* culture was tied with tie-tie techniques and allowed for vegetative growth along the long line.



Figure 1.8 Floating raft cultivation system at Lombok Island, Indonesia.



Figure 1.9 The *Kappaphycus* seedlings were attached to lines within a floating raft cultivation system.

Generally, there are four types of *Gracilaria* cultivation methods: the open water cultivation, pond farming, polyculture, and tank cultivation (Critchley, 1993).

1.4.3 Open Water Cultivation

Open water cultivation is typically conducted in bays, estuaries, upwelling locations, and coral flats. Bottom stocking and rope farming are the two techniques utilized in open sea cultivation. Bottom stocking is the simplest way for cultivating *Gracilaria*. This approach involves transferring vegetative thalli that are naturally attached to small stones and shells to the desired cultivation site. At the cultivation site, vegetative thalli will be planted with a fork in soft sediments or affixed to the bottom of cultivation sites using sand-filled plastic tubes. This labor-intensive technique harvests *Gracilaria* when the biomass reaches 15 kg m⁻².

Two types of rope farming are available. The first method involves attaching vegetative thalli to a rope and then placing them at the bottom of the cultivation site. The second method of rope cultivation utilizes reproductive elements such as spores. In this procedure, a rope is used to capture the *Gracilaria* spores at their habitat. Then this rope, along with the germlings attached to it, will then be transferred back to the cultivation site for cultivation.

1.4.4 Pond Farming

In China, pond farming is a common practice. Annually, 2000 tonnes of dry *Gracilaria* are harvested from 1500 hectares of ponds (Figure 1.10). Each pond is typically 0.7–1.0 hectares in size and 60–70 cm deep. By regulating the pond water depth, the salinity of the seawater is kept between 10 and 20 ppt, the pH is kept between 7 and 8, and the temperature is kept between 15 and 30 degrees Celsius. In this technique, ponds are planted with 5–6 tonnes per hectare of *Gracilaria* and supplemented with urea (1 gL⁻¹) and other nutrients. 30–50 percent of the entire biomass is typically gathered for every 30–35 days of cultivation. *Gracilaria* is also grown in polyculture systems with economically significant organisms like tilapia, milkfish, prawns, crabs, and abalone.



Figure 1.10 Pond farming cultivation system at Muar, Johor, Malaysia. (Laman Alamjaya Sdn Bhd). Photo credit: Dr Emienour Muzalina Mustafa, UMT.

1.4.5 Tank Farming

Generally, tank farming is a costly type of cultivation. In this technique, the entire process or a portion of the process is automated. As a standard component, an aeration system is crucial for promoting the circulation of seaweed within the tank. In this technique, nutrients like nitrogen-based fertilizer are added. *Gracilaria* is often picked on a frequent basis to maintain a high level of output (Figure 1.11 and 1.12).



Figure 1.11 Selected seaweed tank cultivation system used for the *Gracilaria* culture, at the hatchery, Universiti Malaya.



Figure 1.12 Growth of *Gracilaria* culture in the tank cultivation system.

1.5 Advanced Techniques of Seaweed Cultivation

Since 1990, the seaweed aquaculture industry has significantly increased its production rate, demonstrating 30 years of steady growth and emergence as a viable industry. It has also become one of the key sources of revenue, particularly for coastal communities in several tropical nations, as an alternative means of subsistence. However, over time, some decline in seaweed productivity has been observed in some countries due to the use of vegetative propagation techniques over a long period of time, resulting in strain fatigue, or loss of vigour (Reddy et al. 2017). To address this issue, some efforts are being made to identify new technologies, one of which is the use of micropropagation techniques to produce high-quality seedlings or plant materials for the industry.

In-vitro cell culture technology has facilitated the development of micropropagation in seaweed cultivation since the year 1978. *Chondrus crispus* and *Laminaria angustata* are seaweed species that accomplished tissue culture in 1978 (Reddy et al. 2008). The initial research on this regeneration study is focusing on the preparation of axenic explants with the fundamental aspects of seaweed tissue culture (Chen & Taylor 1978; Reddy et al. 2003; Yokoya & Yoneshigue-Valentin 2011). Seaweed micropropagation, including protoplasm culture, spore culture, tissue culture, and somatic embryogenesis (Prabowo et al. 2021), is selected to produce disease-free and healthy callus tissues for mass proliferation. Polne-Fuller and Gibor (1987) reported the earliest callus induction by tissue culture studies on *Eucheuma* and *Kappaphycus*, followed by Dawes and Koch (1991), Bradley (1991), Reddy et al. (2003), and Yeong et al (2014). These studies comprised the development of suitable sterilizing procedures for explants, the development of callus induction protocols with suitable induction medium containing plant growth regulators, and the improvement of the growth and regeneration of callus culture by enhancing the culture media. Several following investigations have shown the effective regeneration of micro-propagules directly from explants of *E. serra* (Sahoo et al. 2002), *K. striatus* (Yunque et al. 2011), and *K. alvarezii* (Sahoo et al. 2003, Neves et al. 2015; Yong et al. 2014, Hurtado et al. 2015). As reported by Reddy et al. (2003) and Yeong et al. (2014), these tissue and callus culture techniques provide a clonal propagation method for maintenance of seedstock for seaweed aquaculture, a micropropagation technique for selection of high-quality germplasm and improvement of productivity of economically valuable strains, such as *K. alvarezii*.

As compared to the traditional vegetative propagation, tissue/callus culture and associated micropropagation techniques have provided a strategy for improving the quality and quantity of high-value seedstock and planting materials for the seaweed aquaculture industry. To fulfill the rising worldwide demand for biomass, this high-quality seedstock must be supplied to aquaculture farming in a more effective manner. This may be accomplished using photobioreactors for the purpose of enhancing the growth and productivity of *K. alvarezii* under controlled conditions with important abiotic growth factors such as light irradiance, temperature, mixing time, and the supply of plant growth regulators and nutrients (Muoz et al. 2006 and Yong et al. 2014). These reports have been published in scientific journals. The successful implementation of bioreactors not only provides a scaled-up system for mass production of plant materials for a chosen strain, but also allows for use

in overcoming the obstacles of seasonality and environmental variability that hinder the growth of seaweed cultures and in bioprocess engineering production (Rorrer & Cheney, 2004; Reddy et al. 2008 and Yong et al. 2014).

In addition to a monoculture approach, the seaweed aquaculture sector might use integrated multitrophic aquaculture (IMTA) to cultivate seaweed alongside fish, shrimp, or mollusks. This IMTA system enables the cultivation of numerous species, including flora and animals, inside a system in which the engaged species mutually benefit one another. The IMTA may be conducted in either an open system (near-shore or at sea) or an enclosed system. In the IMTA model, the co-products (organic and inorganic wastes) of one fauna species are recycled to serve as nutritional input for the other species, and in exchange, the fauna species in the system offer clean water with oxygen to the other fauna species.

The use of IMTA (open system) near-shore or open seaweed farming between or adjacent to fish, shrimp, or mollusk aquaculture was able to mitigate the negative ecological effects of aquaculture while increasing seaweed yield. The land-based enclosed IMTA system with a recirculation aquaculture system and multiple biosensors, as well as controlled culture conditions, was able to optimize the integration of excreted ammonium, phosphates, and CO² by seaweeds, and accomplish a cleaning environment for fish and shrimp that offered a more balanced nutrient flow within the system. An important benefit of this method is that it allows the primary farmed species to turn their wastes into additional commercial and environmental value by converting them into sub products like fertilizers, food, and energy instead of aquaculture waste, which enables stronger long-term and overall sustainability and profitability of the culture (Gacía et al., 2020).

The concept of developing a sustainable seaweed cultivation technique is to minimize the negative impact of aquaculture on the environment and at the same time, maximize the product derived from seaweed cultivation. In particular, the seaweed aquaculture industry emphasizes the recovery of optimum biomass of harvested seaweed and then creating a more sustainable ecosystem. Despite this potential, seaweed cultivation acts as a hub for nutrient removal in coastal waters. Scientists are estimating that dissolved inorganic nitrogen going into the coastal ecosystem will escape the shelf break or increase by 2 times by year 2050 compared to year 1990 (Davis et al. 2003; Jickells et al. 2018; Malone and Newton 2020) with one option for recycling of excessive nutrients being in seaweed cultivation technique. While most cultivation technologies only focus on developing a robust and cost-effective system, it is important to note that the advanced technique of seaweed cultivation must be able to improve the condition of coastal waters.

Nutrient bioextraction with seaweed is one of the approaches that can increase the seaweed biomass without utilizing unsustainable nutrients (Park et al. 2021). Through this approach, seaweed cultivation not only supports the food webs and biomass but is also able to improve the quality of coastal waters (Wu et al. 2017; Xiao et al. 2017). It is generally agreed that biosorption or

bioremediation using seaweed is one of the most eco-friendly mechanisms for balancing nutrient management and mitigating eutrophication (La Barre et al. 2018; Rubín et al. 2006). To increase the efficiency of the nutrient bio-extraction method, IMTA is implemented as a cost-effective cultivation method while performing the synergistic functions of utilizing the organic and inorganic nutrients in the ecosystem. In the era of Industry 4.0, IMTA, or real-time environmental monitoring or biosensors, through engineering mechanisms, can provide in-situ analysis for better seaweed cultivation planning. However, this data is only able to maintain mainly sustainable seaweed cultivation.

Global climate trends have influenced environmental parameters and the distribution of aquatic diseases, contributing to seaweed's low survival rates. Genetic engineering is applied to manipulate the genes involved in stress tolerance and growth rate to increase the quality and quantity of seaweed. Many investigations have demonstrated the advantages of using molecular tools to study the adaptation of tropical seaweed in fluctuating environmental conditions. Gene expressions of seaweed involved in environmental stresses were studied by Nikaido et al. (2000) with *Porphyra yezoensis* using a large expressed sequence tag (EST). After that, different molecular tools such as cDNA microarray, real-time polymerase chain reaction (qPCR) and transcriptomic were used to investigate the defense pathway related to seaweed stress responses, including *Gracilaria changii*, *Kappaphycus alvarezii*, *Eucheuma cottoniil* and *Sargassum binderi* (Baharum et al. 2011. Ho et al. 2008; Teo et al. 2009, Smolina et al. 2016; Liesner et al. 2020). Molecular genetic information provides seaweed's environmental tolerance limits and vulnerability to abiotic stresses. With the information provided from the above molecular tools, the molecular priming technique can be applied to generate stress memory in seaweed. This technique is more acceptable than genetically modified organisms (GMOs) and can ensure the production security of seaweed under environmental stresses. However, priming for cultivation still needs to overcome a challenge as mitotically grown generations will reset their genetic memories during meiosis and embryogenesis.

The success of seaweed cultivation will be influenced by internal factors and environmental conditions. Clearly, the factors contributing to enhancing the defense mechanism and level of stress tolerance of seaweed must be investigated through genetic engineering and genetic manipulation approaches. In conjunction with genetic engineering to improve the seaweed strains, multi-disciplinary techniques for seaweed cultivation must be designed to help intensive cultivation of higher biomass seaweed and increase sustainability.

1.6 Conclusion

Since seaweed can produce promising raw materials for food, animal feed, medicine (including nutraceutical and pharmaceutical products), energy (biofuel; biofuel cell), and other uses, there is a need to increase current seaweed production or expand into new seaweed varieties through the development of seaweed cultivation technologies. For the seaweed aquaculture industry, it is essential to maintain research efforts aimed at the development of high-quality seed stocks with properties of disease resistance, rapid growth rate, and a high concentration of desired biochemical

compounds, which can be accomplished through micropropagation and bioprocess engineering. New research efforts are also needed in the enhancement of seaweed cultivation techniques, especially with the integration of mechanization technologies for more efficient and eco-friendly aquaculture systems and efficient post-harvest technologies, for the continued growth of the seaweed aquaculture industry. It also ensures the seaweed aquaculture industry's long-term stability and sustainability.

References

- Abdul Khalil, H. P. S., Tye, Y., Saurabh, C., Peng, L. C., Lai, T. K., Chong, E., Fazita, M., Jaafar, M. H., Banerjee, A. and Syakir, M. I., 2017. Biodegradable polymer films from seaweed polysaccharides: A review on cellulose as a reinforcement material. *Express Polymer Letters*, 11, 244-265.
- Abdullah Al-Dhabi, N. and Arasu, M., 2022. Biosorption of hazardous waste from the municipal wastewater by marine algal biomass. *Environmental Research*, 204, 112115.
- Baharum, H., Morita, H., Tomitsuka, A., Lee, F. C., Ng, K. Y., Rahim, R. A., Abe, I., & Ho, C. L. (2011). Molecular cloning, modeling, and site-directed mutagenesis of type III polyketide synthase from *Sargassum binderi* (Phaeophyta). *Marine biotechnology* (New York, N.Y.), 13(5), 845–856. <https://doi.org/10.1007/s10126-010-9344-5>
- Baweja P., Kumar S., Sahoo D., Levine I. (2016). Biology of Seaweeds In *Seaweed in Health and Disease Prevention*, Fleurence J & Levine I. (Eds) (pp. 41-106) Academic Press, Elsevier. United Kingdom.
- Bixler, H. J., & Porse, H. (2010). A decade of change in the seaweed hydrocolloids industry. *Journal of Applied Phycology*, DOI 10.1007/s10811-010-9529-3
- Bradley P (1991) Plant hormones do have a role in controlling growth and development of algae. *J Phycol* 27:317–321
- Buhani, Wijayanti, T., Suharso, Sumadi and Ansori, M., 2021. Application of modified green algae *Nannochloropsis sp.* as adsorbent in the simultaneous adsorption of Methylene Blue and Cu(II) cations in solution. *Sustainable Environment Research*, 31 (17).
- Cai, J., Lovatelli, A., Aguilar-Manjarrez, J., Cornish, L., Dabbadie, L., Desrochers, A., Diffey, S., Garrido Gamarro, E., Geehan, J., Hurtado, A., Lucente, D., Mair, G., Miao, W., Potin, P., Przybyla, C., Reantaso, M., Roubach, R., Tauati, M. & Yuan, X. 2021. Seaweeds and microalgae: an overview for unlocking their potential in global aquaculture development. FAO Fisheries and Aquaculture Circular No. 1229. Rome, FAO
- Chen, L.C.M. and Taylor, A.R.A. (1978). Modularity tissue culture of the red alga *Chondrus crispus*. *Can. J. Bot.* 56: 883-6. In Baweja, P., Sahoo, D., Garcia-Jimenez, P., & Robaina, R.R. (2009). Review: Seaweed tissue culture as applied to biotechnology: Problems, achievements and prospects. *Phycological Research*, 57.
- Cherry, P., O'Hara, C., Magee, P., McSorley, E. and Allsopp, P., 2019. Risks and benefits of consuming edible seaweeds. *Nutrition Reviews*, 77 (5), 307-329.
- Davis, T.A., Volesky, B. and Mucci A. (2003). A review of the biochemistry of heavy metal biosorption by brown algae. *Water research*. 37(18):4311-4330. DOI: [10.1016/s0043-1354\(03\)00293-8](https://doi.org/10.1016/s0043-1354(03)00293-8)
- Dawes C, Koch E (1991) Branch, micropropagule and tissue culture of the red alga *Eucheuma denticulatum* and *Kappaphycus alvarezii* farmed in the Philippines. *J Appl Phycol* 3:247–257
- Eka Putri, L., Syafiq, E. and Apriliani, E., 2018. Adsorption Efficiency of Lead by *Sargassum crassifolium* in Different Biomass Size and Dose. *Proceedings of the 1st International Conference on Recent Innovations*, 1809-1814.
- FAO., 2022a. Fishery and Aquaculture Statistics. Global aquaculture production 1950-2020 (FishStatJ). In: FAO Fisheries and Aquaculture Division [online]. Rome. Updated 2022. www.fao.org/fishery/statistics/software/fishstatj/en
- FAO. 2022b. Fishery & Aquaculture Statistics. Global production by production source 1950-2020 (FishStatJ). In: FAO Fisheries and Aquaculture Division [online]. Rome. Updated 2022. www.fao.org/fishery/statistics/software/fishstatj/en
- Gacia, S.; Leandro, A.; Cotas, C.; Cotas, J.; Marques, J.C.; Pereira, L.; Gonçalves, A.M.M. (2020) The evolution road of seaweed aquaculture: Cultivation technologies and the industry 4.0. *Int. J. Environ. Res. Public Health*, 17, 6528.
- Ganesan, A., Tiwari, U. and Rajauria, G., 2019. Seaweed nutraceuticals and their therapeutic role in disease prevention. *Food Science and Human Wellness*, 8 (3), 252-263.
- García-Poza, S.; Leandro, A.; Cotas, C.; Cotas, J.; Marques, J.C.; Pereira, L.; Gonçalves, A.M.M. (2020). The Evolution Road of Seaweed Aquaculture: Cultivation Technologies and the Industry 4.0. *Int. J. Environ. Res. Public Health*, 17, 6528.
- Ho, CL., Teoh, S., Teo, SS. et al. Profiling the Transcriptome of *Gracilaria changii* (Rhodophyta) in Response to Light Deprivation (2009). *Mar Biotechnol* 11, 513–519 . <https://doi.org/10.1007/s10126-008-9166-x>
- Hurtado A. Q. and Critchley A. T. (2006) In: *Advances in Seaweed Cultivation and Utilization in Asia*, Phang, S. M, A. Crichery, Ang P. (eds) (pp. 21-28). University of Malaya Maritime Research Centre (UMMRc), University Malaya, Kuala Lumpur.
- Hurtado AQ, Neish IC, Critchley AT (2015) Developments in production technology of *Kappaphycus* in the Philippines: more than four decades of farming. *J Appl Phycol* 27:1945–1961
- Isam, M., Baloo, L., Kutty, S. and Yavari, S., 2019. Optimisation and Modelling of Pb (II) and Cu (II) Biosorption onto Red Algae (*Gracilaria changii*) by Using Response Surface Methodology. *Water*, 11 (11), 2325.
- Jensen, A. (1993). Present and future needs for algae and algal products. *Hydrobiologia*, 260/261, 15-23.

- Jickells, T. D., et al. (2017), A reevaluation of the magnitude and impacts of anthropogenic atmospheric nitrogen inputs on the ocean, *Global Biogeochem. Cycles*, 31, 289– 305, doi:10.1002/2016GB005586.
- Kasim, M., Balubi, A.M., Astuti, O., Rahman, A., Patadjai, R.S., Muskita, W.H., Takwir, A., Ruslaini, Bahtiar, & Jalil, W. (2021). Comparison between the growth of *Kappaphycus alvarezii* (Rhodophyta) seed from tissue culture and clone selection cultivated using horizontal net. *The Egyptian Journal of Aquatic Research*.
- Khalid, S., Abbas, M., Saeed, F., Bader-UI-Ain, H. and Rasul Suleria, H. A., 2018. Therapeutic Potential of Seaweed Bioactive Compounds. *Seaweed Biomaterials*, 1-12.
- Kulal, P. and Badalamoole, V., 2020. Hybrid nanocomposite of kappa-carrageenan and magnetite as adsorbent material for water purification. *International Journal of Biological Macromolecules*, 165, 542-553.
- La Barre, S., S. Bates, S., Neveux, N., Bolton, J.J., Bruhn, A., Roberts, D.A. and Ras, M. (2018). The Bioremediation Potential of Seaweeds: Recycling Nitrogen, Phosphorus, and Other Waste Products. In *Blue Biotechnology* (eds S. La Barre and S. S. Bates). <https://doi.org/10.1002/9783527801718.ch7>
- Liesner, D., Fouqueau, L., Valero, M., Roleda, M. Y., Pearson, G. A., Bischof, K., Valentin, K., & Bartsch, I. (2020). Heat stress responses and population genetics of the kelp *Laminaria digitata* (Phaeophyceae) across latitudes reveal differentiation among North Atlantic populations. *Ecology and Evolution*, 10(17), 9144–9177. <https://doi.org/10.1002/ece3.6569>
- Lindsey White W. & Wilson P. (2015) World seaweed utilization In: *Seaweed Sustainability*, Brijesh K. Tiwari, Declan J. Troy (Eds) (pp.7-25). Academic Press Elsevier. United Kingdom.
- Lohroff, T. J., Gillette, P. R., Close, H. G., Benetti, D. D., & Stieglitz, J. D. (2021). Evaluating the potential bioextractive capacity of South Florida native macroalgae *Agardhiella subulata* for use in integrated multi-trophic aquaculture (IMTA). *Aquaculture*, 544, [737091]. <https://doi.org/10.1016/j.aquaculture.2021.737091>
- Mahdavinia, G. and Etemadi, H., 2019. Surface modification of iron oxide nanoparticles with κ-carrageenan/carboxymethyl chitosan for effective adsorption of bovine serum albumin. *Arabian Journal Of Chemistry*, 12(8), 3692-3703.
- Malone, TC and Newton, A (2020): *The Globalization of Cultural Eutrophication in the Coastal Ocean: Causes and Consequences*. *Front. Mar. Sci.* 7:670. DOI: 10.3389/fmars.2020.0067
- McHugh, D.J. (2002). Prospects for seaweeds production in developing countries. FAO Fisheries Circular. No. 968. Rome, FAO. p28.
- Moey, S., Abdullah, A. and Ahmad, I., 2015. Edible films from seaweed (*Kappaphycus alvarezii*). *International Food Research Journal*, 22(6), 2230-2236.
- Mouritsen Ole G. (2013) *Seaweeds: Edible, Available & Sustainable* (pp 1-225) University of Chicago Press, Chicago & London.
- Muñoz J, Cahue-López AC, Patiño R, Robledo D (2006) Use of plant growth regulators in micro-propagation of *Kappaphycus alvarezii* (Doty) in airlift bioreactors. *J Appl Phycol* 18:209–218
- Nessim, R., Bassiouny, A., Zaki, H., Moawad, M. and Kandeel, K., 2011. Biosorption of lead and cadmium using marine algae. *Chemistry and Ecology*, 27 (6), 579-594.
- Neves FAS, Simioni C, Bouzon ZL, Hayashi L (2015) Effects of spindle inhibitors and phyto regulators on the micro-propagation of *Kappaphycus alvarezii* (Rhodophyta, Gigartinales). *J Appl Phycol* 27:437–445
- Nikaido I, Asamizu E, Nakajima M, Nakamura Y, Saga N, Tabata S. 2000. Generation of 10,154 expressed sequence tags from a leafy gametophyte of a marine red alga, *Porphyra yezoensis*. *DNA Research* 7: 223–227.
- Oualid, H., Abdellaoui, Y., Laabd, M., El Ouardi, M., Brahmi, Y., Iazza, M. and Oualid, J., 2020. Eco-Efficient Green Seaweed *Codium decortatum* Biosorbent for Textile Dyes: Characterization, Mechanism, Recyclability, and RSM Optimization. *American Chemical Society*, 5 (35), 22192–22207.
- Park, J-S, Shin, SK, Wu, H, Yarish, C, Yoo, HI, Kim, JK. (2021). Evaluation of nutrient bioextraction by seaweed and shellfish aquaculture in Korea. *J World Aquac Soc.* 52: 1118– 1134. <https://doi.org/10.1111/jwas.12786>
- Pereira, R & Yarish, C. (2008). Mass Production of Marine Macroalgae. In: *Ecological Engineering*. Vol.3 of Encyclopedia of Ecology, 5 vols. (pp. 2236-2247) Oxford: Elsevier.
- Phang S.M., Shaharuddin S., Noraishah H. & Sasekumar A. (1996). Studies on *Gracilaria changii* (Gracilariales, Rhodophyta) from Malaysian mangroves. *Hydrobiologia*, 327: 347-352. DOI: 10.1007/BF00047829
- Phang S.M. (2010). Potential products from tropical algae and seaweeds especially with reference to Malaysia. *Malaysian J. Sci.*, 29(2): 160-166.
- Phang S.M., Lim P.E. & Yeong H.Y. (2010a). Malaysian seaweed resources in the South China Sea and their potential economic and ecological applications. *J. Sci. & Tech. in the Tropics*, 6(2): 87-109.
- Phang S.M., Yeong H.Y., Lim P.E., Md Nor A.R. & Gan K.T. (2010b). Commercial varieties of *Kappaphycus* and *Euclima* in Malaysia. *Malaysian J. Sci.*, 29(3): 214-224.
- Prabowo, B.H., Kurnianto, D., Aprilla, I.R. and Amilia, S. (2021). The Development and Potential of Seaweed Tissue Culture. *Indonesian Journal of Biology Education*. 4(20): 7-13.
- Reddy C.R.K, Gupta V., Jha B. (2010) Developments in biotechnology of red algae. In: Seckbach J, Chapman DJ (eds) *Red algae in the genomic age, cellular origin, life in extreme habitats and astrobiology* 13. Springer, Dordrecht, pp 307–341
- Reddy C.R.K, Jha B, Fuhita Y, Ohno M (2008) Seaweed micro-propagation techniques and their potentials: an overview. *J Appl Phycol* 20:609–617
- Reddy, C. R. K., Yokoya, N. S., Yong, W. T. L., Luhan, M. R. J., & Hurtado, A. Q. (2017). Micro-propagation of *Kappaphycus* and *Euclima*: Trends and prospects. In A. Q. Hurtado, A. T. Critchley, & I. C. Neish (Eds.), *Tropical Seaweed Farming Trends, Problems and Opportunities* (pp. 91-110).
- Reddy, C.R.K., Jha, B., Fujita, Y. et al. (2008) Seaweed micropropagation techniques and their potentials: an overview. *J Appl Phycol* 20, 609–617.

- Reddy, C.R.K., Kumar, G.R.K., Siddhanta, A.K., Tewari, A. and Eswaran, K. (2003), In Vitro Somatic Embryogenesis And Regeneration Of Somatic Embryos From Pigmented Callus Of *Kappaphycus Alvarezii* (Doty) Doty (Rhodophyta, Gigartinales). *Journal of Phycology*, 39: 610-616.
- Reddy, C.R.K., Yokoya, N.S., Yong, W.T.L., Luhan, M.R.J., Hurtado, A.Q. (2017). Micro-propagation of *Kappaphycus* and *Eucheuma*: Trends and Prospects. In: Hurtado, A., Critchley, A., Neish, I. (eds) *Tropical Seaweed Farming Trends, Problems and Opportunities*. *Developments in Applied Phycology*, vol 9. Springer, Cham. https://doi.org/10.1007/978-3-319-63498-2_5
- Renn, D. (1997). Biotechnology and the red seaweed polysaccharide industry: status, needs and prospects. *Trend in Biotechnology*, 15, 9-14.
- Rorrer GL, Cheney DP (2004) Bioprocess engineering of cell and tissue cultures for marine seaweeds. *Aquac Eng* 32:11–41
- Rubín, E., Rodríguez, P., Herrero, R. and Sastre de Vicente, M.E. (2006), Biosorption of phenolic compounds by the brown alga *Sargassum muticum*. *J. Chem. Technol. Biotechnol.*, 81: 1093-1099. <https://doi.org/10.1002/jctb.1430>
- Sahoo D, Ohno M, Hiraoka M (2002) Laboratory, field and deep seawater culture of *Eucheuma serra* – a high lectin yielding red alga. *Algae* 17:127–133
- Salgueiro, A., Daniel-da-Silva, A., Girão, A., Pinheiro, P. and Trindade, T., 2013. Unusual dye adsorption behavior of carrageenan coated superparamagnetic nanoparticles. *Chemical Engineering Journal*, 229, 276-284.
- Sartal CG, Alonso MC, Barrera PB. 2012. Application of seaweeds in the food industry. In: *Handbook of marine macroalgae: biotechnology and applied phycology*, Kim SK (Ed) (pp. 522–531). John Wiley & Sons, NJ.
- Shah D, K., Brahmabhatt, N. and Thaker, P., 2021. Study of Green Seaweed Biochar for Lead Adsorption from Aqueous Solution. *Oriental Journal Of Chemistry*, 37 (5), 1242-1247.
- Smolina I, Kollias S, Jueterbock A, Coyer JA, Hoarau G. 2016 Variation in thermal stress response in two populations of the brown seaweed, *Fucus distichus*, from the Arctic and subarctic intertidal. *R. Soc. open sci.* 3: 150429. <http://dx.doi.org/10.1098/rsos.150429>
- Soares, S., Simões, T., Trindade, T. and Daniel-da-Silva, A., 2017. Highly Efficient Removal of Dye from Water Using Magnetic Carrageenan/Silica Hybrid Nano-adsorbents. *Water, Air, & Soil Pollution*, 228(3), 87.
- Tahiluddin, A. B. and Terzi, E. (2021). Ice-ice disease in commercially cultivated seaweeds *Kappaphycus* spp. and *Eucheuma* spp.: A review on the causes, occurrence, and control measures. *Marine Science and Technology Bulletin*, 10(3), 234-243. <https://doi.org/10.33714/masteb.917788>
- Teo S-S, Ho C-L, Teoh S, Raha AR, Phang SM. 2009 Transcriptomic Analysis of *Gracilaria changii* (Rhodophyta) in Response to Hyper- and Hypoosmotic Stresses. *J Phycol* 45(5):1093–1099.
- Tirtawijaya, G., Negara, B.F.S.P., Lee, J. H., Cho, M. G., Kim, H.K., Choi, Y. S., Lee, S. H., Choi, J. S. (2022) The Influence of Abiotic Factors on the Induction of Seaweed Callus. *J. Mar. Sci. Eng.* 10, 513. <https://doi.org/10.3390/jmse10040513>
- Turan, G.; Neori, A. (2010) Intensive seaweed aquaculture: A potent solution against global warming. In *Seaweeds and Their Role in Globally Changing Environments*; Israel, A., Einav, R., Seckbach, J., Eds.; Springer: Dordrecht, The Netherlands, pp. 357–372.
- Wan, H., Li, C., Mahmud, S. and Liu, H., 2021. Kappa carrageenan reduced-stabilized colloidal silver nanoparticles for the degradation of toxic azo compounds. *Colloids and Surfaces A: Physicochemical and Engineering Aspects*, 616, 126325.
- Ward, G.M., Kambey, C.S., Faisan, J.P., Tan, P., Daumich, C., Matoju, I., Stentiford, G.D., Bass, D., Lim, P., Brodie, J., & Poong, S. (2021). Ice-Ice disease: An environmentally and microbiologically driven syndrome in tropical seaweed aquaculture. *Reviews in Aquaculture*.
- Wu, H., Kim, J. K., Huo, Y., Zhang, J., & He, P. (2017). Nutrient removal ability of seaweeds on *Pyropia yezoensis* aquaculture rafts in China's radial sandbanks. *Aquatic Botany*, 137, 72-79. <https://doi.org/10.1016/j.aquabot.2016.11.011>
- Xiao, X., Agusti, S., Lin, F. et al. (2017). Nutrient removal from Chinese coastal waters by large-scale seaweed aquaculture. *Sci Rep* 7, 46613. <https://doi.org/10.1038/srep46613>
- Yeong HY, Phang SM, Reddy CRK, Khalid N (2014) Production of clonal planting materials from *Gracilaria changii* and *Kappaphycus alvarezii* through tissue culture and airlift photobioreactors. *J Appl Phycol* 26:729–746
- Yew, Y., Shameli, K., Miyake, M., Kuwano, N., Bt Ahmad Khairudin, N., Bt Mohamad, S. and Lee, K., 2016. Green Synthesis of Magnetite (Fe₃O₄) Nanoparticles Using Seaweed (*Kappaphycus alvarezii*) Extract. *Nanoscale Research Letters*, 11 (1), 276.
- Yokoya, N. , and Yoneshigue-Valentin, Y. (2011). Micropropagation as a tool for sustainable utilization and conservation of populations of Rhodophyta. *Revista Brasileira de Farmacognosia*, 21 (2). doi: 10.1590/s0102-695x2011005000077
- Yong WTL, Ting SH, Yong YS, Thien VY, Wong SH, Chin WL, Rodrigues KF, Anton A (2014) Optimization of culture conditions for the direct regeneration of *Kappaphycus alvarezii* (Rhodophyta, Solieriaceae). *J Appl Phycol* 26:1597–1606
- Yunque DA, Tibubos KR, Hurtado AQ, Critchley AT (2011) Optimization of culture conditions for tissue culture production of young plantlets of carrageenophyte *Kappaphycus*. *J Appl Phycol* 23:433–438

Chapter 2

ADVANCES IN SHRIMP AQUACULTURE TECHNOLOGY

Key-Seem Phang¹, Qi-En Pung¹ and Jiun-Yan Loh^{1,2}*

¹ Faculty of Applied Sciences, UCSI University, 56000 Kuala Lumpur, Malaysia.

² Centre of Research for Advanced Aquaculture (CORAA), UCSI University, 56000, Cheras, Kuala Lumpur, Malaysia

*Corresponding author: lohjy@ucsiuniversity.edu.my

ABSTRACT

Scientific applications, technology, and innovations have accelerated the growth of shrimp aquaculture in the past few decades to meet the ever-increasing global protein demands of the expanding human population. Despite challenges and issues associated with worsening environmental conditions, high energy consumption, disease outbreaks, and production losses, the shrimp aquaculture industry is integrating to maximize production by applying shrimp culture technologies that promote the efficient use of time and energy, security, traceability, and sustainability. This article discusses the technologies and applications that have the right properties to address the mentioned challenges, including disease prevention measures and water recirculation from the applied aspects of biotechnology.

KEYWORDS: Shrimp farming, Sustainable aquaculture, Technology, Innovation

2.1 Introduction

Shrimp aquaculture products represent the largest animal protein industry in the consumer market. As crustaceans are nutritious and provide high number of certain nutrients such as iodine that aren't abundant in many other foods, crustaceans are critical in people's lives (Li & Xiang, 2013). According to the Food and Agriculture Organization (FAO) of the United Nations, the world's shrimp production was approximately 8.63 million tons in 2018 and is predicted to increase by 7% (Shinn *et al.*, 2018). The total shrimp production from aquaculture and capture fisheries is assumed to contribute heavily to the global food supply as the human population grows. The world's shrimp production will exceed

the poultry and pork world's production in the coming years. Despite the relatively poor knowledge of the cultivated shrimp species and the enormous loss due to disease outbreaks, an estimated approximately 1 billion dollars would be generated every year from the global shrimp consumer market (Li & Xiang, 2013). However, there are many problems and challenges such as disease outbreaks caused by the *Vibrio* pathogen, deterioration of environmental parameters, outdated production systems, and deteriorated water parameters, which could harass the development of shrimp aquaculture and its production. These issues are the main limiting factors in shrimp aquaculture and have resulted in serious economic losses.

Technology and science have benefited almost every aspect of human life in the past decades. The introduction of these technologies and its broadened knowledge over the past years, such as utilizing live feed as aquafeed, disease management technology, including microalgae and zooplankton in hatcheries, have also resolved the bottleneck issues in culturing certain marine species and able to promote the rapid development of aquaculture (Conceição *et al.*, 2010). Although these innovative technologies have contributed to the incredible growth of shrimp aquaculture, the challenges shrimp aquaculture faces are daunting in meeting the expanding demand of the consumer market. For more sustainable and profitable development of shrimp aquaculture, a systemic improvement of existing technologies and the implementation of new technology and innovative feeding methods must make crustaceans' production become more effective, economical, environmentally sustainable, and compliant with governmental regulations. These emerging innovative technologies, including information and automation technologies, recirculating aquaculture systems (RAS), molecular technologies for genetic improvement, biofloc technology, etc., are paving the way toward greater productivity and profitability.

2.2 Common Diseases in Shrimp Farming

Shrimp disease has been a major setback to the shrimp farming industry since the 1980s, when the industry was expanding, and the shrimps were routinely shipped from one region to another. Bacterial diseases caused by *Vibrio* spp. are the major cause of the high mortality rate (up to 90%) in shrimp hatcheries, particularly during early larval stages (Li & Xiang, 2013). Diseases such as Taura Syndrome (TS), White Spot Disease (WSD), and Acute Hepatopancreatic Necrosis Diseases (AHPND) could result in production collapse and cost the shrimp industry billions of dollars in loss in terms of jobs and export revenue (Lightner, 2011). The diseases caused by these pathogens and viruses have been far more problematic to manage as shrimp and the damaging pathogens share the same aquatic environment. The emergence of these shrimp diseases mentioned above results from the complex interactions among the host, the environmental variables, and the surrounding microflora (Xiong *et al.*, 2016). Furthermore, the international movement of live and dead aquaculture products has allowed the transfer of pathogens and diseases from country to country and from one continent to another before the etiology is understood. These diseases in the penaeid shrimp industry could also threaten humans as pathogens and viruses can cross the species barrier (Lightner, 2011).

2.3 Disease Prevention and Management

Due to the lack of adaptive immunity systems in shrimps, classic vaccinations against diseases do not work for them. Although the application of antibiotics could alleviate bacterial diseases, it is not effective against viral diseases. Hence, novel technologies are urgently needed to develop an effective strategy for shrimp disease control.

2.3.1 Vaccination

The immunization of aquaculture started decades ago, and vaccination is an effective method to prevent bacterial and viral diseases. However, as traditional vaccination require each individual to be injected manually, it is considered a labor-intensive process that could cause unnecessary stress to the aquaculture species. Ideally, good vaccine should be safe for the shrimps and provide the host with long-lasting protection (Sivasankar *et al.*, 2017). Hence, immersion and oral administered vaccines as alternative vaccination methods have been studied and developed as they are much easier to handle and require minimal handling skills. This minimizes the damage to the shrimp culture and reduces the mortality rate during vaccination. One of the known oral vaccines in the shrimp aquaculture market is AquaVac™Vibromax™, a *Vibrio* inactivated vaccine manufactured by a UK-based company. It has shown promising results in enhanced growth and survival of shrimp at the post-larvae stage (Amatul-Samahah *et al.*, 2020). However, controlling vibriosis by relying solely on vaccination is unlikely and probable. This should be accompanied by proper sanitization and biosecurity measures.

2.4 Functional Feed from Natural Sources as Alternative Protein Source for Aquafeed

The rapid growth of the aquaculture industry and the ever-increasing demand for farmed marine finfish have resulted in spikes in fishmeal prices over the past few years. The commercial production supply of fishmeal has also faced environmental challenges such as the disruption of nutrition profile, nutritional limitations such as imbalanced amino acid profiles, and the risk of immunological competence with the invasion of pathogens, which eventually lead to disease outbreaks. Hence, the incorporation of different nutritional sources from plants, insects, microorganisms, and natural medicinal products that contain a wide variety of bioactive compounds into shrimp feed has been brought into discussion by researchers and scientists. These innovative sources could maintain the growth and physiological performance and simultaneously enhance shrimps' immunity and disease resistance ability without causing drug resistance.

Innovative feed sources of insect meal and natural medicinal products such as guava (*Psidium guajava*) leaf extract, tiger milk mushroom (*Lignosus rhinocerotis*), black soldier fly (*Hermetia illucens*), and yellow mealworm (*Tenebrio molitor*) (Figure 2.1) contain several types of polysaccharides such as antimicrobial peptides (AMP), β and α -glucans, and chitin, which are potential prebiotics and crucial classes of effector antimicrobial agents (Van Doan *et al.*, 2019; Xia *et al.*, 2021). Compared to current commercial antibiotics that target one specific mode of action, the

crude extracts of natural products contain multiple bioactive components that could generate multifaceted responses in bacterial inhibition and are less likely to cause antimicrobial resistance.



Figure 2.1 Leaves of guava (top left); Tiger Milk Mushroom (bottom left); Yellow Mealworms (top right); Black Soldier Flies (bottom right) (Adámková *et al.*, 2017; Alma, 2019; Koe, 2022; Saanvi, 2021)

Single-cell proteins (SCPs) represent alternative feed sources that could be prepared from different microbial sources, including microalgae, yeast, other fungi, and bacteria. SCPs can fulfil the protein needs in the shrimp aquaculture industry as they offer high-quality protein ingredients applicable across species. Feeding trials showed that aquaculture species such as salmon, tilapia, and white-leg shrimp fed with SCPs show improvement in physiological performance (Jones *et al.*, 2020). The study further validated SCP capability to replace fishmeal in aquafeed. Further research regarding the effect of SCPs on pathogenic resistance still needs to be conducted. Despite few publications, these innovative alternative feed sources have great potential to be a nutritious, immune-boost, and sustainable aquafeed that should be tested in the future. These new sources also offer the potential of increasing nutritional value and anti-microbial effects through genetic manipulation (Sánchez-Muros *et al.*, 2014)

2.5 Novel Molecular Technologies for Genetic Improvement

Due to the rapid development of sequencing technologies and the reduction of costs in DNA genotyping, genetic modification or genetic improvement could be the crucial key factors to the rapid development of the shrimp aquaculture industry. The combination of molecular genomic technologies with the traditional breeding method could significantly enhance the genetic improvement of the

shrimp species and their immunity capability by increasing their resistance to bacterial and virus diseases. Genetic improvement through genome editing using clustered regularly interspaced short palindromic repeats (CRISPR/Cas9), and zinc finger nucleases (ZFNs) has shown the ability to increase the physiological performance and immunity parameters of aquaculture species (Yang *et al.*, 2022). The application of genome editing allows a quick introduction of desirable alleles into the genomes, increasing the frequency of desired alleles at the loci by determining important traits, generating new alleles, and introducing favorable alleles from other species (Shen & Yue, 2019). Due to the high fecundity and external fertilization of aquaculture species, shrimps are considered suitable for genome editing of many individuals simultaneously (Gratacap *et al.*, 2019). Furthermore, the advancement of high-density single-nucleotide polymorphism (SNP) arrays and routine genotyping by sequencing the selection of desired genome alleles offers a higher selection accuracy than selection based on phenotypic and pedigree records alone (Gratacap *et al.*, 2019). Advances in genome selection and genome editing are dignified to reshape the shrimp aquaculture industry by improving the economically important traits that enable potentially favorable alleles to be introduced into the shrimp breeding program. Hence, subject to favorable regulatory and consumer perceptions, genome selection, and editing technology offers the potential to significantly improve the sustainable production and immunity enhancement of shrimp aquaculture.

2.6 Advances in Modern Shrimp Aquaculture

Although significant growth has been observed in shrimp aquaculture recently, intensification of shrimp aquaculture comes with the cost of unintended consequences such as disease outbreaks and environmental pollution that hinder the development and sustainability of the shrimp industry. As demonstrated by Lightner (2003), biosecurity is the practice of excluding pathogens from the aquaculture culture systems for disease prevention, and it is crucial to the efforts to ensure the productivity and sustainability of the shrimp aquaculture. In the aquaculture system, water quality plays a major role in the health status of the shrimp. Reports have shown that the sudden changes in water temperature, salinity, dissolved oxygen content, toxicity, and eutrophication level could weaken the innate immunity of shrimp and convert opportunistic pathogens from non-virulent to virulent, leading to poor growth and disease outbreaks (Boutin *et al.*, 2013). With the fluctuating water quality conditions brought about by rainy seasons in Southeast Asian countries, shrimp farmers are shifting to farming models aimed at achieving zero water exchange to keep the shrimp culture environment fully controlled and kept away from disease outbreaks caused by bacteria, viruses, and pathogens. A reduction in water exchange ensures biosecurity by preventing the entry of disease carriers from external sources and promoting environmental sustainability by reducing land and water requirements (Chamberlain, 2005).

2.7 Recirculating Aquaculture Systems

Recirculating Aquaculture System (RAS) has been widely adopted since the 1980s by shrimp farmers, especially those who practice intensive culture to achieve stable water quality parameters. RAS is a technology that involves multi-step treatment of water, including solids removal, filtration,

denitrification, and disinfection to make it suitable for reuse in the culture system of aquatic animals (Figure 2.2). The recirculation system uses biofiltration and oxygenation to improve water quality and productivity.




Figure 2.2 Recirculating Aquaculture System (RAS) within modules described by Taw *et al.*, (2013)

Cultivating shrimp with reduced water exchange is beneficial, especially in certain regions where water is a limited resource (Chamberlain, 2005). RAS is considered one of the most environmentally friendly ways of producing shrimp at a commercially viable level. The benefits of RAS aquaculture in terms of efficiency and sustainability have prompted Charoen Pokphand Foods (CP Foods), the Thai food giant's decision to move all its shrimp farms indoors (Holmyard, 2019). Despite the major issues such as the high investment needed for constructing facilities, logistics and staff upskilling, RAS will be a long-term solution for the shrimp farming industry, enabling better yield, disease control and quality. The large investment budget will have a positive impact on both production and environmental sustainability and eventually outweigh the costs eventually.

2.8 Biofloc Technology

Biofloc technology (BFT) is another efficient and environmentally friendly method to produce shrimp with relatively low operational costs. Biofloc systems maintain water quality by using microbial communities that continuously recycle and reuse nutrients in the environment. Microorganisms are

capable of up-taking and converting wastes in the form of nitrogenous metabolites into microbial proteins that serve as a useful source of dietary proteins for cultured shrimps (Nisar *et al.*, 2022). Originally conceived as a low-cost natural means of clean water, biofloc systems are becoming increasingly popular as they provide an additional carbon source to the aquaculture system, improving feed ingestion and digestion while reducing the feed conversion ratio (Chamberlain, 2005).

Significant infectious disease outbreaks have prompted the implementation of strict biosecurity measures, such as close monitoring of the shrimp culture environment and reducing water exchange. Various findings have proven the ability of the BFT to control unfavorable chemicals and toxic substances in aquaculture systems. Production intensity in BFT systems is revealed to be higher than in non-BFT culture systems (Nisar *et al.*, 2022). Nowadays, creative and inventive shrimp-industry players are creating and implementing BFT in conjunction with other advances to achieve more productive integrated systems that produce more food with less input and resources. Important principles and components from RAS and biofloc technology are being combined to ensure and enhance the scalable efficiency.

More experimentation and research must provide more accurate data regarding the appropriate conditions for different species and carbon sources in the aquaculture system. Shrimp farmers must be trained in the practical application of the BFT to ensure the effective use of biofloc systems, enabling a natural way of producing more seafood sustainably while concurrently improving farm profitability.

2.9 Information/Digital Technologies

The technology of Internet of Things (IoT) allows quick collection and analysis of massive amounts of streaming data across the entire shrimp aquaculture industry. Although IoT is relatively new to the shrimp aquaculture industry, these technologies are proven to bring huge benefits to the sector. With the application of information technology and IoT, the environmental conditions in shrimp breeding sites could be efficiently observed in real-time and with higher coverage by incorporating underwater cameras and sensors across multiple breeding farms. Furthermore, IoT allows real-time collection and analysis of environmental and water parameters conditions and provides contextual information to ensure the quality of the breeding cage (Jothiswaran *et al.*, 2020). The combination of IoT (Figure 2.3) and robotics can revolutionize shrimp aquaculture in many aspects as it would make the most labor-intensive and risky tasks such as feeding, pond, and net cleaning, behavior monitoring, and sick fish removal be done much more efficiently at a rather low cost (Kamaruidzaman & Rahmat, 2020). The application of automation technology could make the shrimp industry more profitable as robotics can operate continuously without any interruption under difficult conditions and without the need for human assistance (Yue & Shen, 2021). Certainly, introducing automation and information technology would bring substantial changes to the shrimp aquaculture industry. These technologies must be properly applied to the field in the future.




Figure 2.3 The emerging information/digital technologies, including robotics, drones, sensors, artificial intelligence (AI), etc. are applications that increase the shrimp aquaculture production. These technologies are connected with shrimp farms through satellites, the internet of things (IoT), and mobile phones.

2.10 Conclusion

Shrimp aquaculture plays a crucial role in supplying good quality protein to the high-demand consumer market. The requirement for shrimp supplies is expected to increase substantially in the coming years due to the ever-expanding human population. Cutting-edge and novel technologies must address the main challenges such as high production costs and high disease incidences with low survival rates. In the past few decades, the shrimp industry has encountered major setbacks from disease outbreaks, as well as water and land resource challenges. As the industry has grown, so too have inventions and technological advances. Technologies and methods such as innovative feed sources, genome editing, information technologies, recirculating aquaculture systems, and biofloc technology have shown potential in revolutionizing the shrimp industry by offering more sustainable, predictable, and profitable aquaculture. The industry is consolidating, and shrimp farmers are integrating technologies, demonstrating continuous improvement in shrimp farming techniques. With the right approaches and management to apply the available resources effectively, the shrimp industry can achieve a secure and consistent shrimp supply chain.

References

- Adámková, A., Adámek, M., Mlček, J., Borkovcová, M., Bednářová, M., Kouřimská, L., Skácel, J. and Vítová, E., 2017. Welfare of the mealworm (*Tenebrio molitor*) breeding with regard to nutrition value and food safety. *Potravinarstvo Slovak Journal of Food Sciences*.
- Alma Buelva., 2019, Malaysian firms eye black soldier fly pet food protein, viewed 11 May 2022, <https://www.petfoodindustry.com/articles/8769-malaysian-firms-eye-black-soldier-fly-pet-food-protein>
- Alexandratos, N. and Bruinsma, J., FAO ,Food and Agriculture Organization of the United Nations; 2012. *World agriculture towards, 2030*, p.2050.
- Amatul-Samahah, M.A., Omar, W.H.H.W., Ikhsan, N.F.M., Azmai, M.N.A., Zamri-Saad, M. and Ina-Salwany, M.Y., 2020. Vaccination trials against vibriosis in shrimp: A review. *Aquaculture Reports*, 18, p.100471.
- Boutin, S., Bernatchez, L., Audet, C. and Derôme, N., 2013. Network analysis highlights complex interactions between pathogen, host and commensal microbiota. *PLoS one*, 8(12), p.e84772.

- Chamberlain, G., 2005. 'History of shrimp farming', in V. Alday-Sanz, *The shrimp book*, Nottingham University Press, Chicago, pp. 1-34.
- Conceição, L.E., Yúfera, M., Makridis, P., Morais, S. and Dinis, M.T., 2010. Live feeds for early stages of fish rearing. *Aquaculture Research*, 41(5), pp.613-640.
- Gratacap, R.L., Wargelius, A., Edvardsen, R.B. and Houston, R.D., 2019. Potential of genome editing to improve aquaculture breeding and production. *Trends in Genetics*, 35(9), pp.672-684.
- Gudding, R. and Van Muiswinkel, W.B., 2013. A history of fish vaccination: science-based disease prevention in aquaculture. *Fish & shellfish immunology*, 35(6), pp.1683-1688.
- Han, D., Shan, X., Zhang, W., Chen, Y., Wang, Q., Li, Z., Zhang, G., Xu, P., Li, J., Xie, S. and Mai, K., 2018. A revisit to fishmeal usage and associated consequences in Chinese aquaculture. *Reviews in Aquaculture*, 10(2), pp.493-507.
- Holmyard, N., 2019. The inside story: CP Foods move to total shrimp RAS, viewed 3 May 2022, <https://thefishsite.com/articles/the-inside-story-cp-foods-move-to-total-shrimp-ras-1>
- Johnson, K.N., van Hulten, M.C. and Barnes, A.C., 2008. "Vaccination" of shrimp against viral pathogens: phenomenology and underlying mechanisms. *Vaccine*, 26(38), pp.4885-4892.
- Jones, S.W., Karpol, A., Friedman, S., Maru, B.T., and Tracy, B.P., 2020. Recent advances in single-cell protein use as a feed ingredient in aquaculture. *Current opinion in biotechnology*, 61, pp.189-197.
- Jothiswaran, V.V., Velumani, T. and Jayaraman, R., 2020. Application of artificial intelligence in fisheries and aquaculture. *Biotica Research Today*, 2(6), pp.499-502.
- Kamaruidzaman, N.S. and Rahmat, S.N., 2020, May. Water monitoring system embedded with internet of things (IoT) device: a review. In *IOP Conference Series: Earth and Environmental Science* (Vol. 498, No. 1, p. 012068). IOP Publishing.
- Koe Ting Min., 2022. Ready to roar: Malaysia's tiger milk mushroom maker targets overseas growth amid domestic popularity, Viewed at 5 May 2022, <https://www.nutraingredients-asia.com/Article/2022/03/31/Tiger-milk-mushroom-firm-eyeing-expansion-amid-domestic-popularity>
- Li, F. and Xiang, J., 2013. Recent advances in researches on the innate immunity of shrimp in China. *Developmental & Comparative Immunology*, 39(1-2), pp.11-26.
- Lightner, D., 2007. Biosecurity in Shrimp Farming: Pathogen Exclusion through Use of SPF Stock and Routine Surveillance. *Journal of the World Aquaculture Society*, 36(3), pp.229-248.
- Lightner, D.V., 2003. Exclusion of specific pathogens for disease control in a penaeid shrimp biosecurity program. *The World Aquaculture Society*, pp. 81-116
- Nisar, U., Peng, D., Mu, Y. and Sun, Y., 2022. A solution for sustainable utilization of aquaculture waste: a comprehensive review of biofloc technology and aquamimicry. *Frontiers in Nutrition*, 8.
- Saanvi., 2021, Top 10 Guava Leaves Benefits For Skin, Hair & Health, viewed at 5 may 2022, <https://stylesatlife.com/articles/guava-leaves-benefits/>
- Shen, Y. and Yue, G., 2019. Current status of research on aquaculture genetics and genomics-information from ISGA 2018. *Aquaculture and Fisheries*, 4(2), pp.43-47.
- Shinn, A.P., Pratoomyot, J., Griffiths, D., Trong, T.Q., Vu, N.T., Jiravanichpaisal, P. and Briggs, M., 2018. Asian shrimp production and the economic costs of disease. *Asian Fisheries Science* S, 31, pp.29-58.
- Sivasankar, P., John, K.R., George, M.R., Anushalini, S.V., Kaviarasu, D. and Petchimuthu, M., 2017. Prophylactics in shrimp aquaculture health management: A review. *Journal of Entomology and Zoology Studies*, 5(4), pp.1049-1055.
- Taw, N., Saleh, U. and Slamet, B., 2013. Malaysia shrimp project scales up for production in biosecure biofloc modules. *Global Aquaculture Advocate January/ February 2013*, pp. 40-42.
- Van Doan, H., Hoseinifar, S.H., Esteban, M.Á., Dadar, M. and Thu, T.T.N., 2019. Mushrooms, seaweed, and their derivatives as functional feed additives for aquaculture: an updated view. *Studies in Natural Products Chemistry*, 62, pp.41-90.
- Xia, J., Ge, C. and Yao, H., 2021. Antimicrobial peptides from black soldier fly (*Hermetia illucens*) as potential antimicrobial factors representing an alternative to antibiotics in livestock farming. *Animals*, 11(7), p.1937.
- Xiong, J., Dai, W. and Li, C., 2016. Advances, challenges, and directions in shrimp disease control: the guidelines from an ecological perspective. *Applied microbiology and biotechnology*, 100(16), pp.6947-6954.
- Yang, Z., Yu, Y., Tay, Y.X. and Yue, G.H., 2022. Genome editing and its applications in genetic improvement in aquaculture. *Reviews in Aquaculture*, 14(1), pp.178-191.
- Yue, K. and Shen, Y., 2021. An overview of disruptive technologies for farming.

Chapter 3

COPEPODS AS LIVE FEED FOR MARINE LARVICULTURE: ADVANTAGES, CHALLENGES AND FUTURE PERSPECTIVES

Li-Lee, Chew^{1,2} and Ving-Ching, Chong³*

¹ Centre of Research for Advanced Aquaculture (CORAA), UCSI University, 56000, Cheras, Kuala Lumpur, Malaysia

² Faculty of Applied Sciences, UCSI University, 56000 Cheras, Kuala Lumpur, Malaysia

³ Institute of Biological Sciences, University Malaya 50603 Kuala Lumpur, Malaysia

*Corresponding author: chewll@ucsiuniversity.edu.my

ABSTRACT

With the current trend in human population growth and overexploitation of wild fish, aquaculture is the best solution to support the declining fish production worldwide. Although the overall fish production from aquaculture has exceeded the wild fish production in the last few years, it is still unlikely that the marine finfish from aquaculture can surpass the wild production soon. The success of seed production in mariculture is highly dependent on the type of feed given particularly at the first feeding stage of marine fish larvae. While inert feeds progressively improved in nutritional composition, live feeds are still preferred for marine fish larvae rearing since they are natural food to fish larvae. Aquaculture live feeds consists of microalgae, *Artemia*, cladocerans, copepods and rotifers. Among these, copepods constitute the most ideal feed that meet almost all nutritional requirements of marine fish larvae. High nutritional composition (e.g. DHA, free amino acids, pigments and vitamins), swimming behavior and size of copepods are advantages that allow optimal growth of fish larvae. No enrichment is required for copepods, unlike *Artemia* and rotifers with some nutrient deficiencies. Unstable production and expensive microalgal feeds are major constraints that hinder copepods as feeds in large scale aquaculture system. There has been a progressive improvement in copepod culture at commercial scale including the establishment of diapause egg

product and alternative feeds of bacteria and ciliates which can be potentially more cost-effective than microalgae in copepod culture. However, more works on innovation and optimization are still needed if commercialization of copepod live feeds is intended. This chapter aims to review and discuss the biological features, advantages, challenges and future perspectives of copepods as live feeds in marine larviculture. The status and potential of copepod culture based on the Malaysian scenario are also highlighted.

KEYWORDS: Copepods, fish larvae, live feeds, mariculture, nutritional composition

3.1 Introduction

With the increased pressure on global food supply due to the exponential growth of world's human population and its direct and indirect impacts on environment, climate and political status, food security is now the priority of almost all country's policy. The sufficient food supply would ensure the political stability and sustainable development of a country. Therefore, the novel technology is crucial to improve the quality and yield of agricultural products as to meet the pace of human population growth. Agriculture not only involves the production of terrestrial crops and poultry but also farming of aquatic organisms or commonly known as aquaculture. In comparison with other agricultural sectors, aquaculture has experienced the most rapid growth of production in the last two decades. To date, more than half of the global seafood production come from aquaculture and this contribution is expected to increase further in future (FAO, 2020). Even it is unlikely that the production of marine farmed fish can exceed the production of capture fisheries in the near future, mariculture is still the best solution to ease the existing pressure on the overexploited wild fish in ocean.

It is more challenging to culture marine fishes as compared to their freshwater counterparts due to the complex environmental set up and nutritional requirements especially for young larval stages. The first larval feeding is the most critical stage that would ensure the success of seed production in aquaculture. With the incomplete development of digestive system, the specific food criteria are required for optimal growth and survival of fish larvae (Infante & Cahu, 2001; Yúfera & Darias 2007; Hu et al., 2018). As the natural food in the wild, the planktonic organisms such as microalgae, *Artemia*, ciliates, cladocerans, copepods and rotifers are commonly used as live feeds for marine larviculture (Pan et al. 2022). In most cases, the unsuccessful seed production of marine fish is mainly attributed to unstable live feed supply during the critical stage of first larval feeding (Santosh et al., 2018). Several attempts have been made to replace live feeds with inert feeds but the results were not as good as live feeds for marine larviculture (Chen et al., 2006). The inert feeds are generally poor in palatability and quality. This is not suitable for fish larvae with incomplete development of digestive system (Infante & Cahu, 2001; Yúfera & Darias, 2007). Furthermore, the inert feeds tend to aggregate and sink to the bottom, making the feeds not only less accessible to the fish larvae but the uneaten food would also deteriorate the water quality of the rearing tank. As opposed to inert feeds, the planktonic live

feeds are more visible and available to the larvae since they drift in the water column. The movement of planktonic live feeds can also stimulate the hunting of fish larvae as a visual predator and thus initiate feeding of the fish larvae (Conceição et al., 2010; Nielsen et al., 2017; Kandathil et al., 2020).

In order to meet the demand of large-scale larval rearing, live feed must have some basic advantages such as short life cycle, high reproductive rate, hardy, nutritious, optimal size and unique swimming behaviour (Drillet et al., 2011). The common live feeds cultured for marine fish larvae rearing include *Artemia*, copepods and rotifers. Among these animals, copepods constitute the most ideal feed to marine fish larvae since they are the natural prey to most of the fish larvae. This can be supported by the fact that more than 90% of the larval fish diets is predominated by copepods even though there are diverse groups of zooplankton available in marine ecosystems (Sampey et al., 2007; Llopiz, 2013; Quah et al., 2022). The brackish copepod species *Pseudodiaptomus annandalei* was consumed by many juvenile fishes in tropical mangrove estuaries before the ontogenetic shift to macrobenthic diets (Chew et al., 2012). In view of its significance to small fish diets, *P. annandalei* was isolated and cultured as live feed for grouper larvae (Liao et al., 2001; Chen et al., 2006).

Although the efficacy of copepods as live feeds to marine fish larvae has been well acknowledged, the stable and reliable stock of copepods is a key challenge in aquaculture industry. There have been about 60 copepod species successfully grown in captivity since 1960s and the first large scale production was established in 1980s for the euryhaline species *Acartia tonsa* in Denmark. Since then, several dominant species in temperate waters (e.g. *Acartia* spp., *Centropages* spp., *Eurytemora* spp. and *Oithona* spp.) have also been raised (Drillet et al., 2011). In recent years, a total of nine common copepod species from the families Acartiidae, Euterpinidae, Oithonidae, Paracalanidae, Pseudodiaptomidae and Temoridae have been successfully cultured and used for marine larval fish rearing in India (Santosh et al., 2018). Despite the progressing success over time in copepod culture techniques, the production of reliable copepod culture on large scale remains the main challenge (Dhont et al., 2012). This chapter aims to review and discuss the biological features, advantages, challenges and future perspectives of copepods as live feeds in larviculture of marine fish larvae. The status and potential of copepod culture based on the Malaysian scenario are also highlighted.

3.2 Why Copepods?

In comparisons with other animal live feeds such as *Artemia* and rotifers, copepods constitute the most reliable feed to marine fish larvae in terms of nutritional value, size and physical behaviour (Drillet et al., 2011; Santhanam et al., 2019; Burbano et al., 2020). As the most abundant metazoan on Earth, copepods are important intermediaries transferring the energy

source from the base of the food webs to top predators. Many studies on the larval fish diets in marine ecosystems show the high preference of fish larvae towards copepods over other zooplankton taxa (tunicates, various meroplankton) as prey (Sampey et al., 2007; Llopiz 2013; Quah et al., 2022). The

plunge of copepod wild stocks as attributed by anthropogenic impacts had significantly impinged the production of capture fisheries in Atlantic waters (Beaugrand et al., 2003). In mariculture, the fish larvae perform better in terms of growth and survival when feeding them with copepods compared with *Artemia* and rotifers (Shields et al., 1999; Hamre et al., 2002; Rajkumar & Vasagam 2006). Like the wild environments, the low supply of copepod live feeds during the critical larval feeding period in mariculture had led to poor development and survival of fish larvae (Støttrup et al., 1998; Hamre et al., 2005; Busch et al. 2011). The high selectivity on copepods over other zooplankton taxa as food by fish larvae is mainly due to the high nutritional value of copepods particularly the polyunsaturated fatty acids (PUFA) that are essential for marine larval fish growth (Shields et al., 1999; Hamre et al., 2005; Busch et al., 2011; Matsui et al., 2021).

Based on biochemical profiling, copepods appear to have higher unsaturated fatty acids (n-3 HUFA) than that of *Artemia* and rotifers (Kanazawa 1993; Reitan et al., 1994; Bell et al., 2003). Although with the moderate lipid contents, high DHA relative to EPA and ARA (>20) in copepods are the key factor that allows the optimal development and growth of marine fish larvae (Bell et al., 2003; van der Meeren et al., 2008). In contrast, the minimal DHA/EPA/ARA ratios required for normal larval fish growth can only be achieved in *Artemia* and rotifer cultures via HUFA enrichment. However, the assimilation of essential fatty acids by the larval fishes from the enriched *Artemia* and rotifers is still less efficient as compared to copepods. HUFA is located in the phospholipid fraction of copepods while the enriched *Artemia* and rotifers are located in the neutral lipid or triacylglycerol fraction (Coutteau & Mourente 1997; Bell et al. 2003). As HUFA in the phospholipid fraction are more accessible and digestible than that in the triacylglycerol fraction, this conforms the nutritional advantages of copepods over other live feeds in marine larviculture (Izquierdo et al., 2000; Gisbert et al., 2005).

Besides HUFA, copepods also consist of higher free amino acids (FAA), astaxanthin and iodine as compared to *Artemia* and rotifers (van der Meeren et al., 2008). The free amino acids are crucial for protein uptake and encourage a better growth performance during the critical first-feeding stage (Fyhn et al., 1993; Rønnestad et al., 2003; Rønnestad & Conceição 2005). The astaxanthin pigments are the source of antioxidants and vitamins while the iodine is the precursor of thyroid hormones. The lack of astaxanthin and iodine during the developmental period would lead to malpigmentation and abnormal growth of marine fish larvae (Shields et al., 1999). With their high nutritional value especially the HUFA, enrichment of copepod live feeds is normally unnecessary, unlike *Artemia* and rotifers with some nutrient deficiencies.

3.3 Copepod Species as Live Feed in Aquaculture

Copepods are the most abundant and diverse metazoans on Earth which consist of 10 orders and can be found in almost all aquatic environments (Razouls et al., 2022). However, not all copepod species are suitable as live feeds as some species are parasitic while others are carnivorous. Most of the cultured species are planktonic and belong to the three orders, namely Calanoida, Cyclopoida

and Harpacticoida. The calanoids constitute the largest group in marine environments. The genera that have been well studied include *Acartia*, *Calanus*, *Centropages*, *Paracalanus*, *Pseudocalanus* and *Temora* (Mauchline 1998). Among these, species from the genera *Acartia*, *Centropages* and *Eurytemora* have been successfully cultured in Europe (Støttrup, 2003). In Asia-Pacific region, the genera that have been successfully raised for larval fish rearing include *Acartia*, *Bestiolina*, *Paracalanus*, *Parvocalanus*, *Pseudodiaptomus* and *Temora* (McKinnon et al., 2003; Santosh et al., 2018). The advantages of some calanoid species as live feeds include their high nutritional value and the diapause eggs that can be stored and revived when desired (Marcus & Murray 2001; Drillet et al., 2007). The major drawback of calanoid culture is its low tolerance to crowding that may lead to low hatching and survival rates (Peck & Holste 2006; Jepsen et al., 2007). For some omnivorous towards carnivorous species such as *Acartia* and *Centropages*, cannibalism among own species is high and therefore prevents high density culture (Ohno & Okamura, 1988).

The cyclopoids cover a broader range of distribution, living not only as benthic and planktonic forms but also as parasites. The non-parasitic cyclopoids are mostly omnivorous or detritivorous that feed on organic particles from their environments. Most of the harpacticoids are benthic and feed on bacteria from the biofilms or organic detritus. The few common planktonic species of harpacticoids are *Euterpina acutifrons*, *Clytemnestra scutellata* and *Microsetella norvegica*. The advantages of cyclopoids and harpacticoids as live feeds in aquaculture include their broad dietary spectrum other than microalgae, high tolerance to poor water quality environments and thus increase the survival rate, ability to co-exist with the invasive organisms such as ciliates, nematodes and rotifers (Cutts, 2002; Rhodes, 2003; Støttrup, 2003), high reproductive rate and resilient to crowding (Støttrup, 2003). The major drawbacks are their lower nutritional value compared to calanoids and benthic behaviour that are less accessible by planktonic fish larvae. Thus, the planktonic species such as *Oithona* spp. and *E. acutifrons* are more ideal for fish larviculture.

3.4 Culture Methods and Challenges

Similar to other live feeds, copepods can be cultured using extensive, semi-intensive and intensive methods. The extensive and semi-intensive methods are dependent on the source directly or indirectly from the natural environments. Copepods are obtained by filtering the natural seawaters through a certain mesh size sieve. The pitfalls of this method are the introduction of composite organisms including pathogens and predators into the culture system. The high-density copepod culture and low reliance on the well-trained workers can be achieved by intensive and automated culture system (Rhodes, 2003; Støttrup, 2003).

The key success of marine larviculture is highly dependent on the quality of feeds especially at the first feeding stage. Low seed production and abnormal development of larval fishes are always associated with their poor dietary contents in HUFA, free amino acids, vitamins and some essential nutrients (Støttrup et al., 1998; Hamre et al., 2005; Busch et al., 2011). Although copepods are generally accepted as the universal feeds to marine larval fishes, it is challenging to culture copepods

at large scale. There is a risk to introduce pathogens, invasive species or predators to larval fish culture if the seawater is directly obtained from natural environments (Støttrup, 2003; Dhont et al., 2013). The most common issue encountered in copepod culture is its tendency to be infested by stalked ciliates if seawater is not properly treated. The coexisting nematodes and rotifers may outcompete copepods in terms of food and space. Meanwhile, the voracious predators such as ctenophores and chaetognaths can prey on both naupliar and adult copepods. The herbivorous copepods are mainly fed with diatoms which require the special medium (e.g. F₂) for proliferation and regular sub-culture to avoid culture stock crash. The use of cheap and readily available feed like flour and yeasts is not advisable since these substances would deteriorate the water quality of culture. The copepod culture itself requires regular water quality maintenance, stock harvesting and sorting. Although some hatcheries have successfully established the large-scale culture, the process from microalgal food production to copepod culture maintenance is expensive and labour intensive (Santosh et al., 2018).

3.5 Future Perspectives

The commercial copepod live feed products are less common in the market as compared to other aquatic live feeds such as microalgae, *Artemia*, cladocerans and rotifers. Almost all copepod live feed products available in the market are either frozen or preserved, and used mainly for ornamental purpose. This is normally expensive and economically infeasible for large scale aquaculture production. Although the copepod diapause eggs have been successfully produced by an aquaculture product company (i.e. CFEED) in recent years, the initiation of copepod population from diapause eggs is still not user friendly as the cysts from *Artemia* and rotifers. In most cases, copepods are used to feed only the critical early larval stages and replaced with *Artemia* and rotifers for the older stages. There has been almost no report on marine larviculture that exclusively relies on copepods as live feed. This scenario is a result of the unstable copepod production for large scale aquaculture system. Therefore, more developments, innovations and optimization of culture techniques are needed to ease the issues of unsustainable production as well as storage and logistics for copepod live feed products.

To achieve the commercial status like *Artemia* and rotifers, the innovative techniques of how to mass produce viable copepod eggs that can be stored and revived when desired, stands a great potential in marine larviculture. The attempts to preserve and prolong shelf-life of copepod eggs (Drillet et al., 2007) can be further researched to optimize and enhance the copepod live feed reliability. Copepods do not produce cysts that can be harvested and dry-packed, unlike *Artemia* and rotifers. Researchers and live feed producers can adopt the similar concept by converting the copepod diapause eggs or egg sacs to the commercial products that can be stored and used like *Artemia* and rotifer cysts.

Since the production of microalgal feeds is expensive and labour intensive, the development of alternative feeds such as bacteria not only can reduce the operational cost but also as a backup when there is crash in microalgal culture. The addition of commercial probiotics was reported to

significantly improve the egg production and hatching success in *A. tonsa* cultures (Drillet et al., 2011), indicating that bacteria can be potentially used in copepod culture. It has been found that rotifers fed phototrophic bacteria that grown in palm oil mill effluent (POME-Bac) have higher PUFA than the population that was cultured with microalgae (Loo et al., 2015). Although there has been no published work on the use of POME-bac as a feed for copepods, the similar research team has successfully maintained the calanoid culture in their hatcheries. As the main palm oil producer in the world, the conversion of POME to bacteria feed for copepod culture constitutes a great potential of aquaculture industry in Malaysia.

Although the importance of copepods in marine larviculture is generally accepted by the local farmers, most of the marine fish farms in the county are still depending on *Artemia* due to its readily available stock in the market. There are few dominant copepod families found in both brackish and marine waters of Malaysia such as Paracalanidae, Pseudodiaptomidae, Oithonidae and Euterpinae (Chew & Chong 2011; 2016). These families constitute the main diets of both larval and juvenile fishes in Malaysian waters (Chew et al., 2012; Quah et al., 2022). However, almost none of them are used as live feeds in aquaculture industry in Malaysia. Most of the copepod culture works in the country were carried out by research institutions or universities. The species involved were from the families Acartiidae and Oithonidae, while the feeds used were conventional microalgal feeds and baker's yeasts (Rajkumar & Rahman 2016; Rasdi et al., 2021). The outputs might be doable at laboratory-scale but more advance techniques are required if commercial scale is intended. Besides Acartiidae and Oithonidae, future research can be extended to other ecologically important families such as Paracalanidae, Pseudodiaptomidae and Euterpinae, while the feeding trials with other alternative feeds (e.g. bacteria and ciliates) can be considered as a replacement or supplement for microalgal feeds. The collaboration between academic researchers and industrial partners is highly recommended to tackle the exact needs and issues of the end users and establish the commercial copepod live feeds that are effective and user friendly to aquaculture industry.

3.6 Conclusion

With both field and laboratory studies, copepods have been proven to be the most optimal food for marine fish larvae. Despite diverse zooplankton groups in marine ecosystems, only a few dominant copepod species (i.e. *Parvocalanus crassirostris*, *Pseudodiaptomus annandalei*, *Oithona* spp. and *Euterpina acutifrons*) predominate the dietary composition of fish larvae. While these species are abundantly found in natural environments, their proliferation in captivity and application on commercial scale remain a challenge. So far, the diapause eggs of *Acartia tonsa* are developed into the commercial product. However, the *Acartia* species may not be a good candidate as the universal food since most of the fish larvae in tropical waters prefer other copepod preys over *Acartia* (Quah et al. 2022). If the same techniques can be adopted and optimized for the more universal species stated above, there would be a great achievement to resolve the issue of copepod live feed supply that have existed for a long time. With the ongoing pace of research and development on copepod

live feeds, the commercialization of copepod live feeds in aquaculture industry will be realized in the near future.

References

- Beaugrand, G., Brander, K. M., Alistair Lindley, J., Souissi, S., & Reid, P. C. (2003). Plankton effect on cod recruitment in the North Sea. *Nature*, 426(6967), 661-664. doi: 10.1038/nature02164
- Bell, J. G., McEvoy, L. A., Estevez, A., Shields, R. J., & Sargent, J. R. (2003). Optimising lipid nutrition in first-feeding flatfish larvae. *Aquaculture*, 227(1-4), 211-220. doi.org/10.1016/S0044-8486(03)00504-0
- Burbano, M. F., Torres, G. A., Prieto, M. J., Gamboa, J. H., & Chapman, F. A. (2020). Increased survival of larval spotted rose snapper *Lutjanus guttatus* (Steindachner, 1869) when fed with the copepod *Cyclopina* sp. and *Artemia* nauplii. *Aquaculture*, 519, 734912. doi.org/10.1016/j.aquaculture.2019.734912
- Busch, K. E. T., Peruzzi, S., Tonning, F., & Falk-Petersen, I. B. (2011). Effect of prey type and size on the growth, survival and pigmentation of cod (*Gadus morhua* L.) larvae. *Aquaculture Nutrition*, 17(2), e595-e603. doi.org/10.1111/j.1365-2095.2010.00800.x
- Chen, Q., Sheng, J., Lin, Q., Gao, Y., & Lv, J. (2006). Effect of salinity on reproduction and survival of the copepod *Pseudodiaptomus annandalei* Sewell, 1919. *Aquaculture*, 258(1-4), 575-582. doi.org/10.1016/j.aquaculture.2006.04.032
- Chew, L. L., & Chong, V. C. (2011). Copepod community structure and abundance in a tropical mangrove estuary, with comparisons to coastal waters. *Hydrobiologia*, 666(1), 127-143. doi.org/10.1007/s10750-010-0092-3
- Chew, L. L., Chong, V. C., Tanaka, K., & Sasekumar, A. (2012). Phytoplankton fuel the energy flow from zooplankton to small nekton in turbid mangrove waters. *Marine Ecology Progress Series*, 469, 7-24. doi.org/10.3354/meps09997
- Chew, L. L., & Chong, V. C. (2016). Response of marine copepods to a changing tropical environment: winners, losers and implications. *PeerJ*, 4, e2052. doi.org/10.7717/peerj.2052
- Conceição, L. E., Yúfera, M., Makridis, P., Morais, S., & Dinis, M. T. (2010). Live feeds for early stages of fish rearing. *Aquaculture research*, 41(5), 613-640. doi: 10.1111/j.1365-2109.2009.02242.x
- Coutteau, P., & Mourente, G. (1997). Lipid classes and their content of n-3 highly unsaturated fatty acids (HUFA) in *Artemia franciscana* after hatching, HUFA-enrichment and subsequent starvation. *Marine Biology*, 130(1), 81-91. doi.org/10.1007/s002270050227
- Dhont, J., Dierckens, K., Støttrup, J., Van Stappen, G., Wille, M., & Sorgeloos, P. (2013). Rotifers, *Artemia* and copepods as live feeds for fish larvae in aquaculture. In *Advances in aquaculture hatchery technology* (pp. 157-202). Woodhead Publishing. doi.org/10.1533/9780857097460.1.157
- Drillet, G., Lindley, L. C., Michels, A., Wilcox, J., & Marcus, N. H. (2007). Improving cold storage of subitaneous eggs of the copepod *Acartia tonsa* Dana from the Gulf of Mexico (Florida–USA). *Aquaculture Research*, 38(5), 457-466. doi.org/10.1111/j.1365-2109.2007.01673.x
- Drillet, G., Frouël, S., Sichlau, M. H., Jepsen, P. M., Højgaard, J. K., Joarder, A. K., & Hansen, B. W. (2011). Status and recommendations on marine copepod cultivation for use as live feed. *Aquaculture*, 315(3-4), 155-166. doi.org/10.1016/j.aquaculture.2011.02.027
- FAO (2020). The State of World Fisheries and Aquaculture 2020. Rome: Food and Agriculture Organization. doi.org/10.4060/ca9229en
- Fyh, H. J., Finn, R. N., Helland, S., Rønnestad, I., & Lømsland, E. R. (2020). Nutritional value of phyto-and zooplankton as live food for marine fish larvae. In *Fish farming technology* (pp. 121-126). CRC Press.
- Gisbert, E., Villeneuve, L., Zambonino-Infante, J. L., Quazuguel, P., & Cahu, C. L. (2005). Dietary phospholipids are more efficient than neutral lipids for long-chain polyunsaturated fatty acid supply in European sea bass *Dicentrarchus labrax* larval development. *Lipids*, 40(6), 609. doi.org/10.1007/s11745-005-1422-0
- Hamre, K., Opstad, I., Espe, M., Solbakken, J., Hemre, G. I., & Pittman, K. (2002). Nutrient composition and metamorphosis success of Atlantic halibut (*Hippoglossus hippoglossus*, L.) larvae fed natural zooplankton or *Artemia*. *Aquaculture Nutrition*, 8(2), 139-148. doi.org/10.1046/j.1365-2095.2002.00201.x
- Hu, J., Liu, Y., Ma, Z., & Qin, J. G. (2018). Feeding and development of warm water marine fish larvae in early life. In *Emerging Issues in Fish Larvae Research* (pp. 275-296). Springer, Cham. doi: 10.1007/978-3-319-73244-2_10
- Infante, J. Z., & Cahu, C. L. (2001). Ontogeny of the gastrointestinal tract of marine fish larvae. *Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology*, 130(4), 477-487. doi: 10.1016/s1532-0456(01)00274-5
- Izquierdo, M. S., Socorro, J., Arantzamendi, L., & Hernández-Cruz, C. M. (2000). Recent advances in lipid nutrition in fish larvae. *Fish Physiology and Biochemistry*, 22(2), 97-107. doi.org/10.1023/A:1007810506259
- Jepsen, P. M., Andersen, N., Holm, T., Jørgensen, A. T., Højgaard, J. K., & Hansen, B. W. (2007). Effects of adult stocking density on egg production and viability in cultures of the calanoid copepod *Acartia tonsa* (Dana). *Aquaculture Research*, 38(7), 764-772. doi.org/10.1111/j.1365-2109.2007.01730.x
- Kandathil Radhakrishnan, D., AkbarAli, I., Schmidt, B. V., John, E. M., Sivanpillai, S., & Thazhakot Vasunambesan, S. (2020). Improvement of nutritional quality of live feed for aquaculture: An overview. *Aquaculture Research*, 51(1), 1-17. doi: 10.1111/are.14357
- Liao, I. C., Su, H. M., & Chang, E. Y. (2001). Techniques in finfish larviculture in Taiwan. *Aquaculture*, 200(1-2), 1-31. doi.org/10.1016/S0044-8486(01)00692-5
- Llopiz, J. K. (2013). Latitudinal and taxonomic patterns in the feeding ecologies of fish larvae: a literature synthesis. *Journal of Marine Systems*, 109, 69-77. doi.org/10.1016/j.jmarsys.2012.05.002

- Loo, P. L., Chong, V. C., Vikineswary, S., & Ibrahim, S. (2016). Waste-grown phototrophic bacterium supports culture of the rotifer, *B. racionius rotundiformis*. *Aquaculture research*, 47(10), 3029-3041. doi.org/10.1111/are.12750
- Marcus, N. H., & Murray, M. (2001). Copepod diapause eggs: a potential source of nauplii for aquaculture. *Aquaculture*, 201(1-2), 107-115. [doi.org/10.1016/S0044-8486\(01\)00514-2](https://doi.org/10.1016/S0044-8486(01)00514-2)
- Matsui, H., Sasaki, T., Kobari, T., Waqalevu, V., Kikuchi, K., Ishikawa, M., & Kotani, T. (2021). DHA Accumulation in the Polar Lipids of the Euryhaline Copepod *Pseudodiaptomus inopinatus* and Its Transfer to Red Sea Bream *Pagrus major* Larvae. *Frontiers in Marine Science*, 14. doi.org/10.3389/fmars.2021.632876
- Mauchline, J. (1998). *Adv. Mar. Biol.* 33: *The biology of calanoid copepods*.
- McKinnon, A. D., Duggan, S., Nichols, P. D., Rimmer, M. A., Semmens, G., & Robino, B. (2003). The potential of tropical paracalanid copepods as live feeds in aquaculture. *Aquaculture*, 223(1-4), 89-106. [doi.org/10.1016/S0044-8486\(03\)00161-3](https://doi.org/10.1016/S0044-8486(03)00161-3)
- Nielsen, R., Nielsen, M., Abate, T. G., Hansen, B. W., Jepsen, P. M., Nielsen, S. L., ... & Buchmann, K. (2017). The importance of live-feed traps—farming marine fish species. *Aquaculture research*, 48(6), 2623-2641. [doi: 10.1111/are.13281](https://doi.org/10.1111/are.13281)
- Ohno, A., & Okamura, Y. (1988). Propagation of the calanoid copepod, *Acartia tsuensis*, in outdoor tanks. *Aquaculture*, 70(1-2), 39-51. [doi.org/10.1016/0044-8486\(88\)90005-1](https://doi.org/10.1016/0044-8486(88)90005-1)
- Pan, Y. J., Dahms, H. U., Hwang, J. S., & Souissi, S. (2022). Recent Trends in Live Feeds for Marine Larviculture: A Mini Review. *Frontiers in Marine Science*, 9. doi.org/10.3389/fmars.2022.864165
- Peck, M. A., & Holste, L. (2006). Effects of salinity, photoperiod and adult stocking density on egg production and egg hatching success in *Acartia tonsa* (Calanoida: Copepoda): optimizing intensive cultures. *Aquaculture*, 255(1-4), 341-350. doi.org/10.1016/j.aquaculture.2005.11.055
- Quah, W. C., Chew, L. L., Chong, V. C., Chu, C., Teoh, C. Y., & Ooi, A. L. (2022). Does structural change in the zooplankton community affect larval fish feeding in anthropogenically disturbed tropical waters?. *Environmental Biology of Fishes*, 105(1), 55-76. doi.org/10.1007/s10641-021-01189-2
- Rajkumar, M. (2006). Suitability of the copepod, *Acartia clausi* as a live feed for Seabass larvae (*Lates calcarifer* Bloch): Compared to traditional live-food organisms with special emphasis on the nutritional value. *Aquaculture*, 261(2), 649-658. doi.org/10.1016/j.aquaculture.2006.08.043
- Rajkumar, M., & Rahman, M. M. (2016). Culture of the calanoid copepod, *Acartia erythroa* and cyclopoid copepod, *Oithona brevicornis* with various microalgal diets. *Sains Malaysiana*, 45(4), 615-620.
- Rasdi, N. W., Ikhwannuddin, M., Syafika, C. A., Azani, N., & Ramli, A. (2021). Effects of using enriched copepod with microalgae on growth, survival, and proximate composition of giant freshwater prawn (*Macrobrachium rosenbergii*). *Iranian Journal of Fisheries Sciences*, 20(4), 986-1003.
- Razouls, C., de Bovée, F., Kouwenberg, J. and Desreumaux, N. 2022. *Diversity and geographic distribution of marine planktonic copepods*. <http://copepodes.obs-banyuls.fr/en>.
- Rhodes, A. (2003). Methods for high density batch culture of *Nitokra lacustris*, a marine harpacticoid copepod. *The big fish bang*. Edited by HI Browman and AB Skiftesvik. Institute of Marine Research, Bergen, Norway, 449-465.
- Rønnestad, I., Tonheim, S. K., Fyhn, H. J., Rojas-Garcia, C. R., Kamisaka, Y., Koven, W., ... & Conceição, L. E. C. (2003). The supply of amino acids during early feeding stages of marine fish larvae: a review of recent findings. *Aquaculture*, 227(1-4), 147-164. [doi.org/10.1016/S0044-8486\(03\)00500-3](https://doi.org/10.1016/S0044-8486(03)00500-3)
- Rønnestad, I., & Conceição, L. E. C. (2005). Aspects of protein and amino acids digestion and utilization by marine fish larvae. *Physiological and ecological adaptations to feeding in vertebrates*, 389-416.
- Sampey A, McKinnon AD, Meekan MG, McCormick MI (2007) Glimpse into guts: overview of the feeding of larvae of tropical shorefishes. *Marine Ecology Progress Series*, 339, 243–257. doi.org/10.3354/meps339243
- Santhanam, P., Jeyaraj, N., Jothiraj, K., Ananth, S., Dinesh Kumar, S., & Pachiappan, P. (2019). Assessing the Efficacy of Marine Copepods as an Alternative First Feed for Larval Production of Tiger Shrimp *Penaeus monodon*. In *Basic and Applied Zooplankton Biology* (pp. 293-303). Springer, Singapore. doi.org/10.1007/978-981-10-7953-5_12
- Santhosh, B., Anil, M. K., Muhammed Anzeer, F., Aneesh, K. S., Abraham, M. V., Gopakumar, G., ... & Unnikrishnan, C. (2018). *Culture techniques of marine copepods*.
- Shields, R. J., Bell, J. G., Luiz, F. S., Gara, B., Bromage, N. R., & Sargent, J. R. (1999). Natural copepods are superior to enriched *Artemia* nauplii as feed for halibut larvae (*Hippoglossus hippoglossus*) in terms of survival, pigmentation and retinal morphology: relation to dietary essential fatty acids. *The Journal of nutrition*, 129(6), 1186-1194. doi.org/10.1093/jn/129.6.1186
- Støttrup, J. G., Shields, R., Gillespie, M., Gara, M. B., Sargent, J. R., Bell, J. G., ... & Fitzgerald, R. (1998). The production and use of copepods in larval rearing of halibut, turbot and cod. *BULLETIN-AQUACULTURE ASSOCIATION OF CANADA*, 41-45.
- van der Meer, T., Olsen, R. E., Hamre, K., & Fyhn, H. J. (2008). Biochemical composition of copepods for evaluation of feed quality in production of juvenile marine fish. *Aquaculture*, 274(2-4), 375-397. doi.org/10.1016/j.aquaculture.2007.11.041
- Yúfera, M., and Darias, M. J. (2007). The onset of exogenous feeding in marine fish larvae. *Aquaculture*, 268, 53–63. [doi: 10.1016/j.aquaculture.2007.04.050](https://doi.org/10.1016/j.aquaculture.2007.04.050)

Chapter 4

THE IMMUNOMODULATORY EFFECTS OF INSECT MEAL IN AQUACULTURE

Yun Liu¹ & Baskaran Gunasekaran^{1,2}*

¹ Faculty of Applied Sciences, UCSI University, 56000 Cheras, Kuala Lumpur, Malaysia.

² Centre of Research for Advanced Aquaculture (CORAA), UCSI University, 56000, Cheras, Kuala Lumpur, Malaysia.

*Corresponding author: baskaran@ucsiuniversity.edu.my

ABSTRACT

Dietary supplement of immunostimulants to maintain fish health and enhance disease resistance has been widely used worldwide. Several insect species have been found to contain a broad range of bioactive substances of optimising fish health. The most studied and applied is Black Soldier Fly (*Hermetia illucens*). The flies are easy to grow and reproduce, full of nutrition, have high feed efficiency and low rearing pollution. It is rich in chitin, antibacterial peptide and lauric acid which adds the potential as an immunostimulant and health promoter of fish culture.

KEYWORDS: black soldier fly, immunostimulant, fish culture

4.1 Introduction

Black soldier fly (BSF) is an important alternative protein source recognized by the Food and Agriculture Organization (FAO) to maintain sustainable economic and environmental development (Nogales et al., 2019). Meanwhile, BSF is rich in bioactive compounds, which can effectively improve the immune state of aquatic animals (Gasco et al., 2021; 2018; Mousavi et al., 2020). There are numerous findings on the immunostimulatory effects of dietary black soldier fly on terrestrial animals. However, the findings in aquaculture are still at an early stage (Mousavi et al., 2020). To study the immune enhancement effect and health level of BSF substitution in various fish species is a hot research question in aquaculture in recent five years.

4.2 Atlantic salmon (*Salmo salar*)

In 2018, the United States approved BSF to be used as food in salmonids (AAFCO, 2019). Few studies about Atlantic salmon have been performed (Li et al. 2021; 2020; 2019). They found that total replacement of fish meal (FM) with 15% inclusion of BSF larvae meal in diet after 8 until 16 weeks did not compromise the gut health of the fish, but to some extent it seemed to affect positively. BSF meal diet induced the same level of infectious pancreatic necrosis virus (IPNV) antibody, immune and barrier-function gene expression profiles when compared to control group with other immunostimulants. Steatosis of enterocytes was observed in the fore and mid intestine in BSF meal diet group as well. BSF group increased the expression of genes related to stress response, immune tolerance and detoxification activity; a higher relative weight of distal intestine was found in fish fed with the BSF meal diet. Meanwhile, BSF fed fish showed less lipid accumulation in enterocytes based on the degree of hyper-vacuolization revealed by histological examination of the pyloric caeca mucosa. BSF supplemented diet also increased microbial richness and diversity of digesta- and mucosa-related gut microbiota. On another hand, Weththasinghe (2021) compared the effects of BSF larvae meal and paste substituting conventional protein sources in Atlantic salmon for 7 weeks on gut health, skin mucus proteome and immune reaction. The results were like the above, showing that BSF had small effects on skin mucus proteome as well as immune reaction. But low to moderate levels of larvae meal with 6.25% and 12.5% or paste with 3.7% and 6.7%, BSF replacement decreased enterocyte steatosis in pyloric caeca. While replacing level up to 25% with larvae meal or 6.7% with paste, BSF improved distal intestine histology. The plasma C-reactive protein linearly decreased with increasing BSF larvae meal level, while the plasma antioxidant capacity linearly increased with increasing BSF larvae paste level.

4.3 Rainbow trout (*Oncorhynchus mykiss*)

For rainbow trout, Huyben (2019), Terova (2019), Rimoldi (2019) studied the effects of partially defatted, fully defatted or full-fat BSF larvae or prepupae meal as substitutes for FM on gut microbiota of the fish for 5 to 12 weeks. The results were almost identical and showed that up to 50% replacement FM with 30% inclusion of BSF had positive effect on gut microbiota of the fish. BSF diets positively modified gut microbial composition, increasing its richness and diversity. For example, after feeding with BSF-based diets, the amount of lactic acid- and butyrate-breeding bacteria increased; the abundance of *Firmicutes* and *Actinobacteria* increased while *Proteobacteria* reduced; the abundance of *Corynebacterium* increased, which was attributed to the production of lipase and the high levels of dietary lipids; the abundance of *Bacillaceae* and *Mycoplasma* significantly increased, which was attributed to the production of chitinase and the high content of dietary chitin. Terova (2020) studied effects of partially defatted BSF larvae meal substituting FM on hepatic methionine (Met) metabolism. The results of 11-week feeding trial indicated that up to 50% of FM replacement with BSF did not negatively affect hepatic Met metabolism in rainbow trout. Particularly, Met availability in the BSF groups directly regulated the transcription levels of two target genes (CBS, cystathionine β synthase which participate in net Met loss; SAHH, S-adenosylhomocysteine hydrolase which is involved in Met resynthesis) in the Met pathway, keeping

an optimal concentration of one-carbon metabolic substrates, i.e., SAM (S-adenosylmethionine): SAH (S-adenosylhomocysteine) ratio in liver tissue. Kumar (2021) found that supplementation of BSF larvae effectively prevented intestinal enteritis induced by soybean meal and using BSF oil is also beneficial to the health of rainbow trout. After 10-week feeding trial, BSF larvae supplementation of 16% down-regulated the prostaglandin and interferon regulatory factor 1 (IRF-1) level in gut, while significantly increased serum lysozyme activity; BSF oil caused no hepatic or intestinal histomorphological change, but tumor necrosis factor, interleukin-8 and IRF-1 in kidney were significantly upregulated along with serum peroxidase activity was significantly increased; the inclusion of bile acid in the BSF oil group significantly upregulated the gene expression of intestinal prostaglandin.

4.4 Bass

There have been reports on feeding BSF meal to largemouth bass (*Micropterus salmoides*), Asian seabass (*Lates calcarifer*), European seabass (*Dicentrarchus labrax*). Several literatures show that appropriate supplementation of BSF meal to the basal diet could improve blood and serum biochemical response, promote health of intestine and liver, enhance the antioxidant activity, decrease oxidative metabolites and down-regulate relative gene expression, up-regulate the immune gene expression, increase disease resistance of bass.

4.5 Largemouth bass (*Micropterus salmoides*)

Fischer (2022) performed an 8-week feeding trial to evaluate effects of full-fat BSF larvae or pre-pupae dietary supplementation in juvenile largemouth bass. It was reported that most of the hepatic/intestinal gene expression in the 11.9% BSF larvae or/and 9.83% BSF pre-pupae treatments generally mirrored those of the fishmeal-based or/and soybean meal-based control diet. The products of these gene expressions are related to growth, lipid metabolism, antioxidant sentinels and inflammatory cytokines, such as insulin-like growth factor (IGF-I) and growth hormone (GH), cholesterol 7 alpha-hydroxylase (CYP7A1) and fatty acid synthase (FASN), superoxide dismutase (SOD), tumor necrosis factor- α (TNF- α) and transforming growth factor- β 1 (TGF- β 1). But intestinal FASN and hepatic TGF- β 1 expression in 9.83% BSF pre-pupae group were down-regulated compared to other groups. However, liver and intestinal histopathology reveal that BSF larvae is better for the fish compared to BSF pre-pupae.

Peng et al. (2021a; 2021b) conducted a 9-week feeding trial to assess dietary BSF pulp on liver and intestine histomorphology, liver lipid metabolism gene expression, antioxidant and immune capacity of largemouth bass. Adding inclusion of 9% BSF pulp in diets had optimal immunostimulatory effects to the fish, considering effect on growth performance meanwhile. Because in this treatment, serum malonaldehyde (MDA) was reduced but acid phosphatase (ACP) was raised, liver peroxisome proliferators-activated receptors α (PPAR α) and lipoprotein lipase (LPL) mRNA down expressed. And in BSF pulp treated diets, intraperitoneal fat deposition was increased, hepatocytes accumulate more lipids accompanied by certain degrees of hyper-vacuolization and inflammatory cell infiltration.

He suggested BSF pulp should be careful to be used in largemouth bass diets. Xu (2021) performed the bacterial challenge experiments with *Aeromonas hydrophila* to largemouth bass fed for 8 weeks. The results indicated that the cumulative mortality rate after 72 h in 1% zymolytic BSF pulp group significantly dropped, while the 5% zymolytic BSF pulp group sharply accelerated. Meanwhile, antioxidant activity increased, and oxidative metabolites reduced in 1% zymolytic BSF pulp group. Furthermore, compared with 0% group, great changes took place in the intestinal microbial composition of 1% group. The dominant phylum Tenericutes changed into Fusobacteria, and the most abundant genus *Cetobacterium* of Fusobacteria replaced *Mycoplasma* of Tenericutes. The results at the level of microbial potential function showed that amino acids, energy and nucleotides metabolism increased, while glycans and lipids metabolism decreased, and α -N-acetylglucosaminidase activity improved notably. Briefly, 1% zymolytic BSF pulp or 9% BSF pulp or 10% BSF pupae or 12% BSF larvae replacing FM for 8-9 weeks was appropriate to largemouth bass. However, it should be noted that excessive addition may cause side effects.

4.6 Asian seabass (*Lates calcarifer*)

Chaklader (2021), Gupta (2020), Hender (2021) were unanimous that BSF diet could be a promising protein source substituting FM in the diets of Asian seabass. But the immunostimulatory effects of dietary BSF on the fish varied quite widely in the above research, the proposed optimal replacement proportion was inconsistent. One of the aspects that contributes to this inconsistency is the feeding formula which was different. Chaklader agreed that FM protein could be totally replaced by the ingredients, containing 30% full-fat or defatted BSF larvae meal (FBSF or DBSF) mixed with 70% poultry by-product meal (PBM). Furthermore, the effect of supplementing protein source with FBSF could be better. After 8 weeks of feeding, comparing with the control group (0BSF-0PBM), all BSF dietary treatment groups performed no significant difference in the serum immune response, 30FBSF-70PBM group showed no obvious hepatic steatosis, a similar skin mucosal barriers response, and no variation of mortality rate infected with *Vibrio harveyi*. It was unnecessary to defat BSF larvae meal to increase production cost.

Meanwhile Gupta does not recommend complete replacement of FM. The adverse impacts on gut health of fish fed without FM but with 18% BSF and 58% varied non-fishmeal ingredients combination was observed after 7 weeks. In Hender's study, feeding trials of BSF protein (22% BSFP) substituting 30% FM, BSF oil (10.5% BSFO) substituting 30% fish oil (FO), BSF protein and BSF oil (22%/10.5% BSFPO) substituting both 30% FM and 30% FO were respectively conducted for 6 weeks. The results revealed that BSFP diet could increase serum bactericidal activity. Neither BSFP nor BSFO had effect on heat shock proteins related to stress (HSP70 and HSP90). While BSFP and BSFPO significantly upregulated expressions of immune-relevant genes such as interleukin-1 β (IL-1 β) and interleukin-10 (IL-10), as well as increased the amount of intestinal and skin mucin cells.

4.7 European seabass (*Dicentrarchus labrax*)

Moutinho (2021) found that lipid peroxidation, superoxide dismutase and catalase activities in the liver of European seabass were decreased after 9 weeks at intermediary BSF inclusion levels, while the dietary inclusion of BSF had no effect on activities of glutathione peroxidase and glutathione reductase. He deemed that 19.5% inclusion of BSF can be adopted for the fish, with no affecting on skin colour, liver health, fillet fatty acid profile and quality, meanwhile may decrease liver and fillet lipid oxidation. Abdel-Latif (2021) conducted a similar feeding trial period but with a little lower BSF inclusion level as Moutinho's study. He found the immunostimulatory effects were more obvious, comparing all BSF groups with the control group. In the aspect of antioxidative capacity, malondialdehyde levels and activities of superoxide dismutase, glutathione peroxidase and catalase were all elevated in a dose-dependent manner. In the aspect of non-specific immunity, serum lysozyme content and activities of phagocytic index and respiratory burst were significantly improved; the gene expression of interleukin-1beta (IL-1 β), interleukin-10 (IL-10) and heat shock protein 70 (HSP70) in liver were upregulated. In the aspect of disease resistance, the relative survival rate challenged with *Vibrio alginolyticus* was significantly increased. He recommended that up to 50% (inclusion of 15%) dietary FM protein could be replaced by BSF larvae meal for European seabass, with the potential efficacy of antioxidative status, immune responses, and disease resistance, and without adverse impact on fish health.

4.8 Siberian sturgeon (*Acipenser baerii*)

Rawski (2021) and Caimi (2020) respectively research the effects of BSF full-fat and defatted larvae meal for FM replacement on liver and intestine health in Siberian sturgeon. Although the feeding trials were very different, the results were essentially in agreement. In Rawski's study with the inclusion of 10% and 15% full-fat BSF in diets for 7 weeks, the portion of pyloric caeca and proximal intestine were increased, the height and area of villi were enhanced. In Caimi 's study with 18.5% and 37.5% BSF diets for 17 weeks, there was no significant effect on histological index in the liver and distal intestine, but the oxidative stress biomarkers have changed at the inclusion level of 37.5%. In the author's opinion, it doesn't matter whether BSF was defatted or not, the point is to focus on the inclusion proportion of BSF in the feeding formula. To avoid adverse impact on the fish health, high level inclusion of BSF in the diet is not recommended.

4.9 Catfish

A 9-week experiment with BSF larvae substituting FM protein was performed to examine the potential alternative effect of BSF in yellow catfish (*Pelteobagrus fulvidraco*) diet in Xiao's research (2018). Among all tested diets, the highest values of serum lysozyme activity was determined in a diet with 25% substitution (inclusion of 11% BSF). Compared with the control group, increased serum lysozyme activity of fish fed with BSF diets was observed in the maximum replacement of 48% FM (inclusion of 22% BSF). Meanwhile, there were no any negative effects on the index of growth performance and feed conversion rate in this group. So, the author recommended that BSF larvae protein replacing up to 48% of FM protein was feasible for yellow catfish with enhanced the

immunocompetence of the fish and without any adverse effects on the growth performance. Hu (2020) conducted a 8-week feeding trial to assess the effects of soybean oil replacement by BSF larvae oil. Although, there was no significant difference in the contents of anti-superoxide anion and malondialdehyde, activities of catalase and superoxide dismutase in the serum of juvenile yellow catfish. Compared with the control group, no significant difference in concentration of tumor necrosis factor- α (TNF- α) was found. When the replacement ratio of BSF oil exceeded 40% (inclusion of 1.6%), the concentrations of anti-inflammatory factor interleukin-10 (IL-10), pro-inflammatory factor interleukin-6 (IL-6) and interleukin-8 (IL-8) in serum of each treatment group were significantly increased, compared with the control group. A 9-week feeding trial was conducted to evaluate the effect of FM replacement with BSF larvae meal on African catfish (*Clarias gariepinus*) fingerlings in Fawole's research (2020). The results are basically like yellow catfish. There was no significant difference in the serum parameters and differential leucocyte counts. The oxidative stress biomarker, malondialdehyde (MDA) content was also similar between BSF based diet and the control, However, the activity of hepatic function enzyme (AST and ALT) was recorded the lowest value in the fish fed with 50% BSF replacement (inclusion of 12%), and the catalase activity improved in this group.

4.10 Zebrafish (*Danio rerio*)

Zebrafish is an important model animal for nutrition research in aquaculture. As an ideal alternative for commercial fish species, it is usually used to preliminarily evaluate the application prospect of novel ingredients before they are applied to aquafeeds (Zarantoniello et al., 2018). There were literatures (Zarantoniello et al., 2018; Lanes et al., 2021) indicated that certain proportion BSF replacing the FM at certain sample time could promote myogenesis-related genes expression. BSF diets could regulate enzymatic hydrolysis activity of chitin. The expression of stress and immune related genes revealed that BSF diets do not impair the fish health. But high BSF substitutions may not play an immunomodulatory role as well as impossible modifications. The above results in zebrafish corresponded to the results in commercial fish species.

4.11 Other species

In the experiment that FM was partially replaced with defatted BSF meal, juvenile clownfish (*Amphiprion ocellaris*, Pomacentridae) did not show significant differences in lipid metabolism and stress response, and intestinal mucosa and liver parenchyma were normal in all experimental groups (Arturo, 2019). Like seabream (*Sparus aurata*), no different indices of poly saturated fatty acids, thrombogenic and peroxidation were observed in the BSF diets groups (Oteri, 2021). Even though the substitution and conclusion proportion in clownfish was high, those in seabream was low, the feeding trial of the two experiments did not overlap at all. After the feeding trial similar with seabream's, immunostimulatory effects were obvious in juvenile large yellow croakers (*Larimichthys crocea*). For example, serum triglyceride (TG) and cholesterol (CHOL) levels significantly decreased with the increase of the replacement level; total antioxidant capacity (T-AOC) of fish liver in the replacing 20% (inclusion of 7%) group and superoxide dismutase activity (SOD) in the replacing 40% (inclusion of 14%) group were the highest among all groups; the lowest liver malondialdehyde (MDA)

content was found in the replacing 40% group (Han, 2020). Neither Lu (2020) nor Couto (2022) recommended complete replacement by BSF in fish diets, whatever substituting for FM or soybean meal (SM). Lu found that BSF could improve the catalase activity of grass carp (*Ctenopharyngodon idellus*) when dietary SM complete replacement (inclusion of 26% BSF). However, when the replacement level was higher than 50% (inclusion of 13% BSF), the villi length and wall thickness in intestine decreased, the abundance of *Shewanella* and *Aeromonas* reduced, comparing with the SM group. In Couto's study, eosinophilic granulocytes and intraepithelial leukocytes numbers were increased in juvenile meagre (*Argyrosomus regius*) fed the 50% replacement BSF diet (inclusion of 30%). In addition, plasma anti-protease activity improved with the increase of dietary BSF level. Summary of the effects of substituting BSF in feed ingredients on health of fish are shown in Table 1.

4.12 Conclusion

The application of BSF in fish feed clearly showed that it can replace FM and other ingredients in fish feed. If a small inclusion of BSF is fed to all fish species, whether it is BSF larvae or pupae, stronger immune activity can be seen. It can significantly improve the antioxidant and anti-inflammatory ability of fish, enhance the immune gene expression and disease resistance, and has intestinal antibacterial activity that improves intestinal health. However, high dosage may have side effects on fish health, and the optimal dosage varies with fish species. Therefore, it is suggested to do more studies before the large-scale applications are performed. In a word, BSF is not only a protein alternative feed material, but also can enhance immune activity and improve health levels.

Table 4.1 Effects of Black soldier fly (*Hermetia illucens*) substituting feed ingredients on health of fish

Fish species	Initial weight	Alternative mode	Recommended proportion		Duration	Immunity	Disease resistance	Antioxidant capacity	Anti-inflammation ability	Intestinal flora	References
			Substitution	Inclusion							
Atlantic salmon (<i>Salmo salar</i>)	34.00 g	A or B replace α	A: 12.5% B: 6.7%	A: 16.13% B: 35.12%	49 d	(+) skin mucus proteome & immune response		(+) plasma antioxidant capacity	(-) enterocyte steatosis in pyloric caeca, (+) distal intestine histology		Weththasinghe et al., 2021
	1.40 kg	A replace α	100%	15%	112 d					(+) microbial richness & diversity in both digesta- & mucosa-associated intestinal microbiota	Li et al., 2021
		A replace α								(+) relative weight of distal intestine	Li et al., 2020
	49.00 g	A replace β	85%	60%	56 d	(+) genes expression of indicative of stress response & immune tolerance			(+) detoxification activity		Li et al., 2019
Rainbow trout (<i>Oncorhynchus mykiss</i>)	201.80 g	A, C or D replace α	45%	30%	35 d					(+) abundance of <i>Firmicute</i> , <i>Actinobacteria</i> , <i>Corynebacterium</i> & <i>Bacillaceae</i> ; (-) <i>Proteobacteria</i>	Huyben et al., 2019

	66.50 g	E replace α	40%	30%	84 d					(+) abundance of <i>Actinobacteria</i> , (-) <i>Proteobacteria</i>	Terova et al., 2019
	66.50 g	E replace α	50%	30%	84 d					(+) bacterial diversity, (-) <i>Proteobacteria</i> , (+) gut abundance of <i>Mycoplasma</i>	Rimoldi et al., 2019
	32 g	A and G respectively replace α and γ	100%	16%	70 d	(+) serum lysozyme & peroxidase activities; (+) kidney IL-8, TNF & IRF1			(-) soybean meal induced intestinal enteritis		Kumar et al., 2021
Largemouth bass (<i>Micropterus salmoides</i>)	4.90 g	Addition of B		<2%	62 d			(-) genes expression of PPAR- α & lipoprotein lipase, (+) Acetyl-CoA carboxylase 1	Hepatocytes filled with lipids with different degrees of vacuolar degeneration & inflammatory cell infiltration		Peng et al., 2021a
				<18%		(+) immune capacity	(+) antioxidant capacity	(+) intraperitoneal fat deposition	Peng et al., 2021b		
	6.02 g	Addition of F		1%	56 d	(+) bactericidal activity	(-) cumulative mortality rates challenged with <i>Aeromonas hydrophila</i>	(+) antioxidant activity, (-) oxidative metabolites			Xu et al., 2021
Asian seabass	2.52 g	Mixture containing A replace α	50%	19%	49 d	(+) innate immune responsive					Gupta et al., 2020

<i>(Lates calcarifer)</i>						genes expression					
	1.74 g	E or/and G replac or/and γ	α 30%	E: 22.1% G: 10.5%	42 d	(+) IL-1 β & IL-10; the number of mucin cells in the gut & skin					Hender et al., 2021
	0.56 g	Mixture containing A or D replace α	100%	A: 3.5% D: 2.8%	56 d	No effects in the serum immune response	Normal proportion of infection rate with <i>Vibrio harveyi</i>				Chaklader et al., 2021
European seabass (<i>Dicentrarchus labrax</i>)	50.00 g	C replace α	45%	19.5%	62 d			Unaffected glutathione peroxidase & glutathione reductase activities			Moutinho et al., 2021
	12.10 g	A replace α	50%	15%	60 d	(+) phagocytic activity, phagocytic index, serum lysozyme & respiratory burst activities; mRNA expression of HSP70), IL-1 β & IL-10 genes	(+) the relative percent of survival challenge with <i>Vibrio alginolyticus</i>	(+) malondialdehyde levels, & catalase, superoxide dismutase, & glutathione peroxidase enzyme activities			Abdel-Latif et al., 2021

Siberian sturgeon (<i>Acipenser baerii</i>)		D replace α	25%	18.5%	118 d			(+) superoxide dismutase activity in mainly kidney, glutathione reductase activity in liver & kidney, catalase activity in both tissues			Caimi et al., 2020
	14.40 g	A replace α and γ	46%	15%	50 d					(+) pyloric caeca & proximal intestine shares, villus height & area	Rawski et al., 2021
Yellow catfish (<i>Pelteobagrus fulvidraco</i>)	2.12 g	G replace δ	100%	4%	56 d					(+) IL-10, IL-6, IL-8 & TNF- α	Hu et al., 2020
	48.50 g	A replace α	48%	22.30%	65 d	(+) values of immune indexes					Xiao et al., 2018
African catfish (<i>Clarias gariepinus</i>)	4.00 g	E replace α	75%	17.20%	60 d			(+) superoxide dismutase & catalase enzyme activities			Fawole et al., 2020
Zebrafish (<i>Danio rerio</i>)		C replace α	25%	10.5%	7, 14, 21 d post-spawning	(+) HSP70, TNF- α & IL-6, chia.1; (-) nr3c1					Zarantoniello et al., 2018
	5.00 mg	D or C replace α	100%	50%	60 d	(+) chia.2, chia.3, & chia.5 genes expression					Lanes et al., 2021

Clownfish (<i>Amphiprion ocellaris</i>)		D replace α	75%	60%	106 d	Stress response gene expressed normally		Lipid metabolism gene expressed normally	Liver parenchyma was normal	Intestinal mucosa was normal	Arturo et al., 2019
Seabream (<i>Sparus aurata</i>)			50%	15.7%				No differences were for poly saturated fatty acids, thrombogenic & peroxidation indices			Oteri et al., 2021
Large yellow croaker (<i>Larimichthys crocea</i>)	50.08 g		40%	14%	49 d			(+) total antioxidant capacity, (-) liver malondialdehyde content			Han et al., 2020
Grass carp (<i>Ctenopharyngodon idellus</i>)	10.10 g	D replace ϵ	50%	13.4%	56 d			(+) catalase activity		(-) the abundance of <i>Aeromonas</i> & <i>Shewanella</i>	Lu et al., 2020
Meagre (<i>Argyrosomus regius</i>)	18.00 g	E replace α	17%	10%	63 d	(+) numbers of eosinophilic granulocytes & intraepithelial leukocytes					Couto et al., 2022

Note: A denotes full-fat BSF larvae meal; B denotes BSF paste; C denotes BSF prepupae; D denotes defatted BSF larvae meal; E denotes partially defatted BSF larvae meal; F denotes zymolytic BSF pulp; G denotes BSF larvae oil. α denotes fish meal; β denotes protein in the diets; γ denotes fish oil; δ denotes soybean oil; ϵ denotes soybean meal.

References

- AAFCO (Association of American Feed Control Officials). (2019). *AAFCO Midyear Meeting Agenda Book*; Hyatt Regency: Savannah, GA, USA; p. 42.
- Abdel-Latif, H. M., Abdel-Tawwab, M., Khalil, R. H., Metwally, A. A., Shakweer, M. S., Ghetas, H. A., & Khallaf, M. A. (2021). Black soldier fly (*Hermetia illucens*) larvae meal in diets of European seabass: Effects on antioxidative capacity, non-specific immunity, transcriptomic responses, and resistance to the challenge with *Vibrio alginolyticus*. *Fish & Shellfish Immunology*, 111, 111-118. <https://doi.org/10.1016/j.fsi.2021.01.013>.
- Arturo, V. A. J., Randazzo, B., Foddai, M., Sanchini, L., Truzzi, C., Giorgini, E., Gasco, L., & Olivotto, I. (2019). Insect meal-based diets for clownfish: Biometric, histological, spectroscopic, biochemical and molecular implications. *Aquaculture*, 498, 1-11. <https://doi.org/10.1016/j.aquaculture.2018.08.018>.
- Caimi, C., Gasco, L., Biasato, I., Malfatto, V., Varello, K., Prearo, M., Pastorino, P., Bona, M. C., Francese, D. R., Schiavone, A., Elia, A. C., Dörr, A. J. M., & Gai, F. (2020). Could dietary black soldier fly meal inclusion affect the liver and intestinal histological traits and the oxidative stress biomarkers of Siberian sturgeon (*Acipenser baerii*) juveniles?. *Animals*, 10(1), 155. <https://doi.org/10.3390/ani10010155>.
- Chaklader, M. R., Howieson, J., & Fotedar, R. (2021). Growth, hepatic health, mucosal barrier status and immunity of juvenile barramundi, *Lates calcarifer* fed poultry by-product meal supplemented with full-fat or defatted *Hermetia illucens* larval meal. *Aquaculture*, 543, 737026. <https://doi.org/10.1016/j.aquaculture.2021.737026>.
- Couto, A., Serra, C. R., Guerreiro, I., Coutinho, F., Castro, C., Rangel, F., Lavrador, A. S., Monteiro, M., Santos, R., Peres, H., Pousão-Ferreira, P., Gasco, L., Gai, F., Oliva-Teles, A., & Enes, P. (2022). Black soldier fly meal effects on meagre health condition: gut morphology, gut microbiota and humoral immune response. *Journal of Insects as Food and Feed*, 1-16. <https://doi.org/10.3920/JIFF2021.0082>.
- Fawole, F. J., Adeoye, A. A., Tihamiyu, L. O., Ajala, K. I., Obadara, S. O., & Ganiyu, I. O. (2020). Substituting fishmeal with *Hermetia illucens* in the diets of African catfish (*Clarias gariepinus*): Effects on growth, nutrient utilization, haematological response, and oxidative stress biomarker. *Aquaculture*, 518, 734849. <https://doi.org/10.1016/j.aquaculture.2019.734849>.
- Fischer, H., Romano, N., Renukdas, N., Kumar, V., & Sinha, A. K. (2022). Comparing black soldier fly (*Hermetia illucens*) larvae versus prepupae in the diets of largemouth bass, *Micropterus salmoides*: Effects on their growth, biochemical composition, histopathology, and gene expression. *Aquaculture*, 546, 737323. <https://doi.org/10.1016/j.aquaculture.2021.737323>.
- Gasco, L., Józefiak, A., & Henry, M. (2021). Beyond the protein concept: Health aspects of using edible insects on animals. *Journal of Insects as Food and Feed*, 7(5), 715-741. <https://doi.org/10.3920/JIFF2020.0077>.
- Gasco, L., Finke, M., & Van Huis, A. (2018). Can diets containing insects promote animal health? *Journal of Insects as Food and Feed*, 4(1), 1-4. <https://doi.org/10.3920/JIFF2018.x001>.
- Gupta, S. K., Fotedar, R., Foysal, M., Priyam, M., Siddik, M. A., Chaklader, M., Dao, T. T. T., & Howieson, J. (2020). Impact of varied combinatorial mixture of non-fishmeal ingredients on growth, metabolism, immunity and gut microbiota of *Lates calcarifer* (Bloch, 1790) fry. *Scientific Reports*, 10(1), 1-13. <https://doi.org/10.1038/s41598-020-72726-9>.
- Han, X. X., Ye, K., Wang, Z. Y., Chen, F. L., Chai, Z. Q., Zhan, W., & Wang, Q. R. (2020). Effect of substitution of fish meal with defatted black soldier fly larvae meal on growth, body composition, serum biochemical parameters, and antioxidant capacity of juvenile large yellow croaker (*Larimichthys crocea*). *Journal of Fishery Sciences of China*, 27(5), 524-535. <https://doi.org/10.3724/SP.J.1118.2020.19306>.
- Hender, A., Siddik, M. A., Howieson, J., & Fotedar, R. (2021). Black soldier fly, *Hermetia illucens* as an alternative to fishmeal protein and fish oil: impact on growth, immune response, mucosal barrier status, and flesh quality of juvenile barramundi, *Lates calcarifer* (Bloch, 1790). *Biology*, 10(6), 505. <https://doi.org/10.3390/biology10060505>.
- Hu, J. R., Yu, C. J., Wang, G. X., Mo, W. Y., & Huang, Y. H. (2020). Effects of dietary soybean oil replaced with black soldier fly larvae oil on growth performance, plasma biochemical indexes and liver lipid droplets of juvenile yellow catfish. *Acta Hydrobiologica Sinica*, 44(4), 717-727. <https://doi.org/10.7541/2020.086>.
- Huyben, D., Vidaković, A., Hallgren, S. W., & Langeland, M. (2019). High-throughput sequencing of gut microbiota in rainbow trout (*Oncorhynchus mykiss*) fed larval and pre-pupae stages of black soldier fly (*Hermetia illucens*). *Aquaculture*, 500, 485-491. <https://doi.org/10.1016/j.aquaculture.2018.10.034>.
- Kumar, V., Fawole, F. J., Romano, N., Hossain, M. S., Labh, S. N., Overturf, K., & Small, B. C. (2021). Insect (black soldier fly, *Hermetia illucens*) meal supplementation prevents the soybean meal-induced intestinal enteritis in rainbow trout and health benefits of using insect oil. *Fish & Shellfish Immunology*, 109, 116-124. <https://doi.org/10.1016/j.fsi.2020.12.008>.
- Lanes, C. F., Pedron, F. A., Bergamin, G. T., Bitencourt, A. L., Dorneles, B. E., Villanova, J. C., Dias, K. C., Riolo, K., Oliva, S., Savastano, D., & Giannetto, A. (2021). Black soldier fly (*Hermetia illucens*) larvae and prepupae defatted meals in diets for zebrafish (*Danio rerio*). *Animals*, 11(3), 720. <https://doi.org/10.3390/ani11030720>.
- Li, Y., Bruni, L., Jaramillo-Torres, A., Gajardo, K., Kortner, T. M., & Krogdahl, Å. (2021). Differential response of digesta- and mucosa-associated intestinal microbiota to dietary insect meal during the seawater phase of Atlantic salmon. *Animal Microbiome*, 3(1), 1-18. <https://doi.org/10.1186/s42523-020-00071-3>.
- Li, Y., Kortner, T. M., Chikwati, E. M., Belghit, I., Lock, E. J., & Krogdahl, Å. (2020). Total replacement of fish meal with black soldier fly (*Hermetia illucens*) larvae meal does not compromise the gut health of Atlantic salmon (*Salmo salar*). *Aquaculture*, 520, 734967. <https://doi.org/10.1016/j.aquaculture.2020.734967>.
- Li, Y., Kortner, T. M., Chikwati, E. M., Munang'andu, H. M., Lock, E. J., & Krogdahl, Å. (2019). Gut health and vaccination response in pre-smolt Atlantic salmon (*Salmo salar*) fed black soldier fly (*Hermetia illucens*) larvae meal. *Fish & Shellfish Immunology*, 86, 1106-1113. <https://doi.org/10.1016/j.fsi.2018.12.057>.

- Lu, R., Chen, Y., Yu, W., Lin, M., Yang, G., Qin, C., Meng, X., Zhang, Y., Ji, H., & Nie, G. (2020). Defatted black soldier fly (*Hermetia illucens*) larvae meal can replace soybean meal in juvenile grass carp (*Ctenopharyngodon idellus*) diets. *Aquaculture Reports*, 18, 100520. <https://doi.org/10.1016/j.aqrep.2020.100520>.
- Mousavi, S., Zahedinezhad, S., & Loh, J. Y. (2020). A review on insect meals in aquaculture: The immunomodulatory and physiological effects. *International Aquatic Research*, 12(2), 100-115. [https://doi.org/10.22034/IAR\(20\).2020.1897402.1033](https://doi.org/10.22034/IAR(20).2020.1897402.1033).
- Moutinho, S., Pedrosa, R., Magalhães, R., Oliva-Teles, A., Parisi, G., & Peres, H. (2021). Black soldier fly (*Hermetia illucens*) pre-pupae larvae meal in diets for European seabass (*Dicentrarchus labrax*) juveniles: Effects on liver oxidative status and fillet quality traits during shelf-life. *Aquaculture*, 533, 736080. <https://doi.org/10.1016/j.aquaculture.2020.736080>.
- Nogales-Mérida, S., Gobbi, P., Józefiak, D., Mazurkiewicz, J., Dudek, K., Rawski, M., Kieronczyk, B., & Józefiak, A. (2019). Insect meals in fish nutrition. *Reviews in Aquaculture*, 11(4), 1080-1103. <https://doi.org/10.1111/raq.12281>.
- Oteri, M., Di Rosa, A. R., Lo Presti, V., Giarratana, F., Toscano, G., & Chiofalo, B. (2021). Black soldier fly larvae meal as alternative to fish meal for aquaculture feed. *Sustainability*, 13(10), 5447. <https://doi.org/10.3390/su13105447>.
- Peng, K., Mo, W., Xiao, H., Wang, G., & Huang, Y. (2021a). Effects of black soldier fly pulp on growth performance, histomorphology and lipid metabolism gene expression of *Micropterus salmoides*. *Aquaculture Reports*, 20, 100737. <https://doi.org/10.1016/j.aqrep.2021.100737>.
- Peng, K., Mo, W., Xiao, H., Hu, J., Zhu, X., Huang, Y., & Wang, G. (2021b). Dietary black soldier fly pulp affects growth, antioxidant and immune capacity of *Micropterus salmoides*. *Journal of Insects as Food and Feed*, 1-8. <https://doi.org/10.3920/JIFF2021.0046>.
- Rawski, M., Mazurkiewicz, J., Kieronczyk, B., & Józefiak, D. (2021). Black soldier fly full-fat larvae meal is more profitable than fish meal and fish oil in Siberian sturgeon farming: The effects on aquaculture sustainability, economy and fish GIT development. *Animals*, 11(3), 604. <https://doi.org/10.3390/ani11030604>.
- Rimoldi, S., Gini, E., Iannini, F., Gasco, L., & Terova, G. (2019). The effects of dietary insect meal from *Hermetia illucens* prepupae on autochthonous gut microbiota of rainbow trout (*Oncorhynchus mykiss*). *Animals*, 9(4), 143. <https://doi.org/10.3390/ani9040143>.
- Terova, G., Ceccotti, C., Ascione, C., Gasco, L., & Rimoldi, S. (2020). Effects of partially defatted *Hermetia illucens* meal in rainbow trout diet on hepatic methionine metabolism. *Animals*, 10(6), 1059. <https://doi.org/10.3390/ani10061059>.
- Terova, G., Rimoldi, S., Ascione, C., Gini, E., Ceccotti, C., & Gasco, L. (2019). Rainbow trout (*Oncorhynchus mykiss*) gut microbiota is modulated by insect meal from *Hermetia illucens* prepupae in the diet. *Reviews in Fish Biology and Fisheries*, 29(2), 465-486. <https://doi.org/10.1007/s11160-019-09558-y>.
- Weththasinghe, P., Lagos, L., Cortés, M., Hansen, J. Ø., & Øverland, M. (2021). Dietary inclusion of black soldier fly (*Hermetia illucens*) larvae meal and paste improved gut health but had minor effects on skin mucus proteome and immune response in Atlantic Salmon (*Salmo Salar*). *Frontiers in Immunology*, 12, 114. <https://doi.org/10.3389/fimmu.2021.599530>.
- Xiao, X., Jin, P., Zheng, L., Cai, M., Yu, Z., Yu, J., & Zhang, J. (2018). Effects of black soldier fly (*Hermetia illucens*) larvae meal protein as a fishmeal replacement on the growth and immune index of yellow catfish (*Pelteobagrus fulvidraco*). *Aquaculture Research*, 49(4), 1569-1577. <https://doi.org/10.1111/are.13611>.
- Xu, F. M., Hou, S. W., Wang, G. X., Gong, J. Y., Zhou, L., Huang, Y. H., Huang X. De., & Liu, L. (2021). Effects of zymolytic black soldier fly (*Hermetia illucens*) pulp as dietary supplementation in largemouth bass (*Micropterus salmoides*). *Aquaculture Reports*, 21, 100823. <https://doi.org/10.1016/j.aqrep.2021.100823>.
- Zarantoniello, M., Bruni, L., Randazzo, B., Vargas, A., Gioacchini, G., Truzzi, C., Annibaldi, A., Riolo, P., Parisi, G., Cardinaletti, G., Tulli, F., & Olivotto, I. (2018). Partial dietary inclusion of *Hermetia illucens* (Black Soldier Fly) full-fat prepupae in zebrafish feed: biometric, histological, biochemical, and molecular implications. *Zebrafish*, 15(5), 519-532. <https://doi.org/10.1089/zeb.2018.1596>.

Chapter 5

LATEST TRENDS IN AQUACULTURE DISEASE MANAGEMENT

Michelle Oi-Yoon Soo^{1,2}, Fook-Choy Yap³*

¹ Centre of Research for Advanced Aquaculture (CORAA), UCSI University, 56000, Cheras, Kuala Lumpur, Malaysia.

² Faculty of Applied Sciences, UCSI University, 56000 Cheras, Kuala Lumpur, Malaysia.

³ Department of Biological Sciences, Faculty of Science, Universiti Tunku Abdul Rahman (UTAR) 31900 Kampar, Perak, Malaysia.

*Corresponding author: michellesoo@ucsiuniversity.edu.my

ABSTRACT

Aquaculture is a thriving economy today to address the high demand of fish consumption by humans and to reduce further depletion of wild fish population. But the demand for a booming aquaculture trade has resulted in multiple problems such as destruction of natural habitats, pollution of water sources and particularly, disease outbreaks in both inland and coastal farms. This article briefly discusses the common diseases in farmed shrimps, fish, bivalves and seaweeds with insights on the latest disease mitigation and management methods in aquaculture farms, ranging from traditional methods to advanced molecular techniques.

KEYWORDS: aquaculture, disease, management, shrimp, fish, bivalve, seaweed

5.1 Introduction

Food security is a major issue today especially with the exponential growth of human population and scarcity of crops due to land depletion and exhausted oceans. Seafood such as fish, shrimps and seaweeds are the highest traded food commodity (FAO, 2012; Stentiford et al., 2012; Naylor et al., 2021) and plays a pivotal role in nutritional and financial security in developing countries (Bene et al., 2015). In Malaysia, aquaculture has emerged as the top 15 global producers of seafood with 521,000 tonnes of total aquaculture production (Fathi et al., 2018). The Kyoto Declaration forecast has unfortunately materialised with global aquaculture becoming increasingly important in terms of supplementing capture fishery production (Jennings et al., 2016). The Bangkok Declaration identified

management of animal health by cooperative action at national, regional, and inter-regional levels as “an urgent requirement for sustaining growth” (FAO, 2000). Good progress has been made in terms of disease identification, diagnostics, treatment, and zone management but recurring issues, for example lice infestations, are still significant barriers to expansion (Rogers et al., 2016). Infectious diseases caused by viruses, bacteria, and protozoans continue to impose yield-limiting effects on aquaculture production. Additionally, the aquatic environment is a dynamic medium capable of transporting pathogens across vast distances (Munro and Gregory, 2009; Murray, 2013). Pathogens can also be accidentally transferred between farms, due to the movement of organisms and vehicles, creating a convoluted network of pathogen transmission (Munro & Gregory 2009; Murray 2013). To control infectious diseases, large quantities of drugs such as antibiotics have been employed freely resulting in the emergence of resistant organisms (Huang et al. 2015; Lafferty et al. 2015). The most basic way to control diseases prior to medication or technology, would be to manage the environmental parameters (i.e., water quality, culture intensity and temperature), pathogen and aquaculture host. A tilt in the balance or alliance of any of these three factors would offset a disease outbreak. However, despite basic farm practices being followed rigidly, disease outbreaks still occur. This review outlines the different disease management trends for common aquaculture species in Malaysia namely shrimps, fish, bivalves, and seaweeds.

5.2 Disease Management in Shrimp Culture

In the 80's, shrimp aquaculture began with farmers capturing broodstock from the wild introducing these wild broodstocks into natural development cycles (Flegel, 2019). The cultivation of juvenile shrimp from postlarval stage used to occur in earthen ponds with water exchange and the use of pelleted feed, resulting in an increase in the number of ponds as well as in stocking density (Flegel, 2019). The first viral outbreak in shrimps occurred in 1990's, with yellow head virus (YHV) in Thailand and white spot syndrome virus (WSSV) in China and Japan. Figure 5.1 depicts the impact of diseases on shrimp production in Thailand for a duration of about 30 years. The figure showed an increase in shrimp production from 1988 to 1995 when the first occurrence of white spot disease (WSD) began, followed by a period of recovery before another episode in a cyclical manner (Thitamadee et al., 2016; Flegel, 2019). The most serious decline in production was due to early mortality syndrome and acute hepatopancreatic necrosis disease (EMS/AHPND), a devastating disease that wipes out entire populations in tanks within hours. This resulted not only in shrimp mortality, but farmers were reluctant to stock in large amounts until they were sure of successful and healthy harvests (Thitamadee et al., 2016).

Open aquaculture systems are subject to occasional disease outbreaks followed by recovery periods. These diseases are sometimes spread by transboundary movement of infected stocks or by uninfected foreign culture stocks being moved to a new area where they are infected with local pathogens (Flegel, 2006; Flegel & Fegan, 2002).

Twenty viruses have been documented to infect penaeid shrimp (Lightner, 1999). Two types of viruses cause deadly outbreaks: DNA viruses, namely the monodon baculovirus (Lightner & Redman, 1981) and the white-spot syndrome virus (WSSV) (Lo et al., 1996), the hepatopancreatic parvovirus and the infectious hypodermal and haematopoietic virus (Lightner, 1996); and RNA viruses, such as the yellow-head virus (YHV), the Taura syndrome virus (Dhar et al, 2004) and the infectious myonecrosis virus (Poulos et al., 2006). WSSV is the most severe threat for farmed shrimp worldwide and is one of the best-studied crustacean viruses today (Walker, & Mohan, 2009; Sanchez-Pas, 2010; Shekar et al., 2012).

Bacterial infections are a major concern in shrimp aquaculture. *Vibrio harveyi* and *V. vulnificus* cause massive larval mortality (Bachere, 2013; Valente & Wan, 2021), whilst *V. parahaemolyticus*, *V. damsela*, *V. nichripulchritudo*, *V. alginolyticus* and *V. penaeicida* cause outbreaks in nurseries or grow-out ponds of shrimps (Bachere, 2013). In 2010, a new disease affecting postlarvae emerged in Asia: a highly virulent strain of *V. parahaemolyticus* that caused the acute hepatopancreatic necrosis disease which acquired a virulent plasmid encoding a pore-forming bacterial toxin (Lee et al., 2015).

The most current methods in mitigating shrimp diseases mostly revolve around DNA technology and biotechnological methods. Farms in Asia are moving towards the production of specific-pathogen free (SPF) stocks that promote faster growth and higher tolerance towards diseases. The introduction of SPF, *Litopenaeus vannamei* in 2000 was so successful that shrimp farmers in Asia only wanted to culture *L. vannamei*. SPF stocks tolerant to TSV were developed using standard breeding and selection methods (Moss et al, 2012). Development of SPF provided farmers with the peace of mind knowing that stocks are 'clean' and not going to cause an outbreak.




Figure 5.1 Shrimp production in Thailand showing the nature of disease outbreaks. AHPND: acute hepatopancreatic necrosis disease; EMS: early mortality syndrome; MSGS: monodon slow growth syndrome; WSD: white spot disease (adapted from Flegel, 2019).

DNA vaccinations were commonly used but the success of the DNA vaccines are based on the antigen–antibody mechanism and are not applicable to shrimp because shrimps don't produce antibodies (Flegel, 2019). The exact mechanism for vaccination is still being studied as it is not fully understood how injection of viral DNA does protect shrimps particularly from white spot disease (WSD) (Chang et al., 2017).

Shrimps produce a large array of antimicrobial peptides (AMP) against bacteria, viruses, and fungi (Bachere et al, 2000; Tassanakajon et al., 2000). It has been found that certain feed ingredients can regulate semi-specific induction (Flegel, 2019). Some AMPs have broad spectrum antibacterial activity, while some may be more effective against gram-negative bacteria. Because of this, feed manufacturers add specific substances to their feed to boost shrimp's immune response (Destoumieux-Garzon et al., 2016). However, it is possible to design an appropriate molecule that could be added to shrimp feed if the shrimp's immune response system is well understood. This approach allows the binding of selected receptors to up-regulate the AMP, which is perfect for defending against pathogens (Flegel, 2019).

The use of dsRNA to protect shrimp against viruses was designed for WSD (Robalino et al., 2005) and for YHD (Yodmuang et al., 2006). This technology was also successful for other shrimp viruses on which it was tried on (Labreuche & Warr, 2013). Using dsRNA as a feed additive is promising if the production cost can be lowered and if the vaccine can be delivered orally (Itsathitphaisarn et al., 2017).

5.3 Disease Management in Fish Culture

With fisheries playing an enormous role in sustaining food security and livelihood, disease outbreaks in aquaculture are resulting in huge losses of production as well as adding constraints to livelihoods and causing food insecurity. China, a major player in aquaculture production suffered a total fish production loss of 15% to disease outbreaks alone (Leong and Bates, 2014). As the saying “prevention is better than cure” goes, preventing the occurrence of diseases go a long way instead of just treating it (Romera et al., 2012). Improved husbandry practices, vaccine, supplements, probiotics/prebiotics, genetically resistant stock, movement restrictions, non-specific immunostimulants, medicinal plant products, water disinfection, biological control, antimicrobial compounds and disinfection of facility water are the some of the approaches in the control of infectious diseases in fish (Kumar et al., 2016).

Diseases in fish aquaculture are multifold and caused by various pathogens ranging from viruses, bacteria, fungi, protozoans and metazoans. Table 5.1 lists down common pathogenic diseases in fish culture farms.

Table 5.1 Common pathogenic diseases in fish culture farms.

Name of disease	Causative Pathogen
Vibriosis	<i>Vibrio</i> spp. including <i>V. anguillarum</i> , <i>V. harveyi</i> clade, <i>V. parahaemolyticus</i> , <i>Aliivibrio salmonicida</i> (<i>V. salmonicida</i>), <i>V. vulnificus</i> , <i>Photobacterium damsela</i>
Edwardsiellosis	<i>Edwardsiella anguillarum</i> , <i>E. ictaluri</i> , <i>E. piscicida</i> , <i>E. tarda</i> , <i>Yersinia ruckeri</i>
Pseudomonas	<i>Pseudomonas anguilliseptica</i> , <i>P. fluorescens</i>
Aeromonas	<i>Aeromonas</i> spp. including <i>Aeromonas caviae</i> , <i>A. hydrophila</i> , <i>A. sobria</i> , <i>A. veronit</i> , <i>A. jandaei</i> , <i>A. salmonicida</i>
Mycobacteriosis	<i>Mycobacterium fortuitum</i> , <i>M. marinum</i> , <i>Nocardia asteroides</i> , <i>N. crassostreae</i> (<i>ostreae</i>), <i>N. seriolae</i>
Streptococcosis	<i>Streptococcus agalactiae</i> , <i>S. iniae</i> , <i>Lactococcus garviede</i> , <i>Aerococcus viridans</i>
Channel catfish virus	Herpesviridae
Infectious Hematopoietic Necrosis	Novirhabdoviridae
Viral Hemorrhagic Septicemia	Novirhabdoviridae
Viral Erythrocytic Necrosis	Iridoviridae
Epizootic Hematopoietic Necrosis	Iridoviridae
Megalocytivirus	Iridoviridae
Viral Nervous Necrosis	Betanodaviridae
Protozoans	<i>Ichthyophthirius multifiliis</i> (ich)
	<i>Trichodinia</i> spp, <i>Trichodonella</i> spp.,
	<i>Ichthyobodo necator</i>
	<i>Apiosoma</i> spp., <i>Ambiphrya</i> spp.
	<i>Chilodonella</i> spp.
	<i>Myxobolus cerebralis</i> (whirling disease)
Metazoans	Isopods
	Monogeneans
	Copepods

Fish can also suffer from non-communicable diseases resulting from poor environmental conditions and malnutrition. For example, the gas bubble disease that can result in gas-filled lumps on the body and eyes due to excessive dissolved gases in the water. Acidosis is also another environmental disease in fish whereby the pH level of the water is too low, resulting in the water becoming too acidic and fish would suffer from acidosis by swimming rapidly and attempting to jump out of the tank. Nutritional deficiencies in fish can cause the following diseases: scoliosis (as shown in Figure 5.2), lordosis, cataract, anemia, fin erosion, stunted growth, gill hyperplasia and reduced feeding, to name a few.



Figure 5.2 Scoliosis and lordosis in a striped mullet, *Mugil cephalus* (adapted from Overstreet & Hawkins, 2017)

A single approach is not enough and cannot sustainably contain a disease as compared to a combination of different strategies which would be more efficient in curbing a disease outbreak in farms. Here, a few approaches to managing diseases along with the most current mitigation methods will be discussed.

Vaccination is an age-old method used to control diseases in aquaculture. Today, there are several variations of vaccination being used in the fish farming scene to manage diseases. Some considerations that should be noted before fish vaccination are the species of fish, production cycle, farming technology, immune status of fish, life history of aquaculture system, the disease in question, the season when the disease occurs, environment, stress factors and costings (Adams, 2016). The types of vaccines used in aquaculture are killed or deactivated vaccines, DNA vaccines, subunit vaccines, genetically-modified vaccines and recombinant vector vaccines (Assefa & Abunna, 2018). Vaccinations are usually done before exposure to a pathogen to allow the fish's immunity to develop. Other factors that determine suitability for vaccination is the size of the fish and temperature of water (Aly & Albutti, 2014). Vaccines can be administered orally, via immersion and via injection.

The use of antibiotics, as prominent as it was back then, is no more a primary treatment option today due to resistance development in fish and an environmental disturbance to the natural microbiota. Due to how disruptive antibiotics can be to both fish and its environment, farmers are advised to use methods apart from antibiotics but if antibiotics must be used due to special circumstances, only approved antibiotics can be administered (Rogers & Basurco, 2009). The American FDA formulated a list of

chemicals/antibiotics for aquaculture that has been reviewed (Assefa & Abunna, 2018) including acetic acid, carbon dioxide, hydrogen, sodium chloride, garlic, onion and ice to reduce the metabolic rate of fish during transport (Bentzon-Tilia et al., 2016). If antibiotics are necessary, it is usually administered orally via medicated feed, injection or by immersion.

Probiotics has emerged as a very popular alternative to antibiotics in recent years due to its many advantages, starting from the improvement of internal gut microbiota. The administration of probiotics can be considered a biological disease control because probiotics contain non-pathogenic bacterial strains that multiply within the fish's intestines to normalise microflora and maintain a microbial balance in the fish (Defoirdt et al., 2011). Probiotics are normally administered through diet, encapsulation and immobilized probiotics by direct admission into water. The more commonly used bacterial strains as probiotics in fish farms are *Bacillus subtilis*, *Lactobacillus acidophilus*, *Lactobacillus sakei*, and *Shewanella putrefaciens*, and these strains have shown great promise in preventing diseases and promoting weight gain in fish (Assefa & Abunna, 2018).

As opposed to probiotics, prebiotics are also a widely used immune booster and are referred to as "food for probiotics" (Assefa & Abunna, 2018). Some commonly used prebiotics are fructooligosaccharide (FOS), mannan-oligosaccharides (MOS), and mixed oligo-dextran (Carbone & Faggio, 2016).

The use of medicinal plants in aquaculture holds promise as plants have been reported to produce various beneficial effects such as anti-stress, growth promotion, appetite stimulation, immune-stimulation, aphrodisiac and to have anti-pathogenic properties in fish aquaculture due to varied active compounds such as alkaloids, terpenoids, tannins, saponins and flavonoids (Reverter et al., 2017). The active components need to be extracted prior to administration orally, via immersion or injection (Reverter et al., 2017).

Biosecurity measures are also an important factor in managing the spread and outbreak of aquatic diseases in fish farms. Farm-level biosecurity measures involve a combination of activities such as strict quarantine measures, equipment sanitation, disinfection of eggs and facilities, traffic control, water parameter management and treatments, clean feed and appropriate disposal of dead/diseased fish (Assefa & Abunna, 2018). These protocols should be enforced during the introduction of new stocks to avoid transferring pathogens (Assefa & Abunna, 2018).

Improving the genetics of fish is also beneficial in disease management. In Singapore, the Temasek Life Sciences Laboratory (TLL) in collaboration with Marine Aquaculture Centre (MAC) started a breeding programme to improve the Asian sea bass's (*Lates calcarifer*) growth performance, disease resistance and meat quality. TLL developed a library of genomic resources consisting of whole genome sequences, DNA markers, a molecular parentage system, linkage maps, BAC libraries, transcriptomes, and genotyping-by-sequencing platform (Shen et al., 2020). After three generations of within-family selection, the Asian sea bass showed a growth increase of 40% and resistance to the big belly disease, nodavirus and iridovirus (Shen et al., 2020).

5.4 Disease Management in Bivalve Culture

Clam, scallop, oyster and mussel farming represents about 50% of the aquaculture market (Zanella et al., 2017). Bivalve aquaculture can be considered a 'cleaner' form of aquaculture due to a self-regulating cultivation mechanism that requires no additional feed, no antibiotic usage and minimal maintenance effort (Tan, 2021). The farming techniques are simple and can be easily transferred to local growers (Tan, 2015). The oyster was a favorite contender of preferred shellfish in the annual food cycle in Malaysia with the production of oysters coming in third after blood cockles, *Anadara granosa* and green mussels, *Perna viridis* (DOF, 2018; Admodisastro et al., 2021).

The major obstacle in the bivalve aquaculture sector is the emergence and spread of infectious diseases that are intensified by high stocking densities (Guo et al., 2016; Pernet et al., 2016; Zanella et al., 2017). Bivalves are filter feeders and due to this habit, these organisms uptake pathogens while feeding. Some common bacteria found in marine water are those from the genus *Vibrio* and because of how abundantly available these bacteria are, they are often a source for disease outbreaks. Various bacteria species, specifically *Vibrio* (*Vibrio alginolyticus*, *V. anguillarum*, *V. tapetis*, *V. tubiashi*, *V. neptunius*, *V. splendidus*, *V. aestuarianus* and other *Vibrio* spp.), *Pseudomonas* and *Aeromonas* (Kueh et al., 1985; Prado et al., 2014; Biel et al., 2014) have been reported to cause bacterial diseases of bivalves which are responsible for bacillary necrosis in the larvae of bivalves (Tubiash et al., 1965; Sugumar et al., 1998). The deciliation, velar epithelial loss and abnormal swimming behaviour are caused by exotoxins (ciliostatic factors and hemolysins) produced by *Vibrio* spp. (Valerio et al., 2010).

The most notable bivalve viruses come from the families Herpesviridae and Iridoviridae (Renault and Novoa, 2004; Zanella et al., 2017). These viral pathogens are highly infectious and transmit easily. The increased disease prevalence in bivalve farms is associated to the high-density systems and environmental changes (Guo and Ford, 2016). The detection of a herpes-related viral infection in oyster was first reported by Farley et al. (1972) in *Crassostrea virginica* from the USA. This oyster herpesvirus infection was later shown to be linked with high death rates, especially in larvae and spawn (Segarra et al., 2010). Recent studies also have shown that the oyster, *C. gigas* in Japan, South Korea, Sweden and Norway was infected by OsHV-1 (Bai et al., 2015; Mortensen et al., 2016), causing mass mortality rate. Ultrastructural analysis revealed that larvae displayed generalized infections, while focal infections occurred mainly in juveniles. Adult stages seemed less prone to infections compared to juveniles. A significant reduction in feeding and swimming activities was observed in the infected larvae, resulting in 100% mortality within days. The infection is usually accompanied with the necrosis of the host cell, specifically the epithelium of velar and mantle. Additionally, it also found to be affecting the swimming activity of the larvae. Histological analysis showed that the abnormal cytoplasmic basophilia and enlarged nuclei with marginated chromatin were observed in the fibroblastic-like cells; other cell types including hemocytes and myocytes showed extensive chromatin condensation (Da Silva et al., 2008).

For iridoviruses, “Maladie des branchies” is the first viral disease recorded on bivalves that was responsible for mass mortalities in the Portuguese oysters, *Crassostrea angulata*, off the French coasts. This gill disease is caused by gill necrosis virus (GNV) where the gills of infected *C. angulata* usually exhibit intense haemocyte infiltration, abscessing and necrosis. “Maladie des branchies” also affects the Pacific oyster, *C. gigas*, imported in France, but the gill lesions re-occurred with a mild mortality. Due to its natural resistance to infection, *C. gigas* has become the main species bred in Europe.

Another type of Irido-like virus known as the “hemocyte infection virus” was reported as the main cause for the mass mortality of *C. angulata* in France between the years of 1970 and 1973. Histopathological assessment revealed that the discolouration of the digestive gland, disruption in connective tissues and hemocytes infiltration (McGladdery, 2011) was observed in the infected oyster. A third Irido-like virus, the oyster velar virus (OVV), caused high mortalities of larval Pacific oysters on the west coast of North America between the years of 1976 to 1984 (Leibovitz et al., 1998). This virus targeted larvae over 150 µm and usually associated with the lesions in the velum and other ciliated epithelia.

Members belonging to the genera *Haplosporidium*, *Bonamia*, *Marteilia* and *Perkinsus* have been identified as the most important protozoan pathogens that target oyster and clam species, causing devastating damage to commercial productions. Diseases caused by *P. marinus*, *P. olseni*, *M. refringens*, *B. ostreae* and *B. exitiosa* are currently under surveillance and require immediate notification by the World Organization for Animal Health. External factors such as temperature and salinity have strong impact on the disease prevalence and intensity of infections (Fernandez Robledo et al., 2014). The pathogen is a lethal infection to the hemocytes, multiplies within blood cells and spreads to all tissues. The pathogen infection is usually accompanied by yellow discoloration in tissues, lesions on gills and mantles, and breakdown of connective tissues. The mortality of bivalves can reach up to more than 90% (Zanella et al., 2017). Apart from the adult, these protozoans can easily infect bivalve larvae. A lack of resistance to infection can easily trigger an epizootic, particularly the high densities of oysters in close-proximity culture beds (Zanella et al., 2017). Perkinsosis is a dermo disease caused by *Perkinsus marinus*, a highly contagious protistan protozoan which is responsible for drastic declines in populations of *C. virginica* (Burreson & Calvo, 1996) and *C. gigas*, *C. rhizophorae* and *C. corteziensis* (Ford, 2011). *Perkinsus olseni* caused perkinsosis in clam populations in Australia, Korea, China, Japan and Europe (Park et al., 2010). Transmission of perkinsosis occurred directly among the oysters; viable cells were released in host feces or the decomposition of the infected host (Park et al., 2010) and were acquired through host feeding mechanisms. This epidemic disease infects every life stage of bivalves (Paynter et al., 2010) whereby the pathogen infects and proliferates in the digestive epithelium, infects the connective tissue of all organs including hemocytes, causing hemocytosis and tissue lysis with severe emaciation (Smolowitz, 2013). *Perkinsus marinus* causes dermo disease which infects the hemocytes of oysters and subsequently affects their growth and fecundity, eventually leading to death. Another common disease in oysters is MSX, or “multinucleated unknown disease” caused by *Haplosporidium nelsoni* which also results in mass mortalities (Figure 5.3).

Research efforts are focused in exploring inexpensive and effective treatments to prevent production losses when diseases occur in hatcheries (Zanella et al., 2017). The “*OIE Manual of Diagnostic Tests for Aquatic Animals*” has detailed protocols to detect specific pathogens that can be used to confirm histological examination results. With such approaches, it can provide an accurate species-specific diagnosis. Conventionally, the disease-causing pathogens are usually assessed using histopathology; however, the histopathological results today is considered a presumptive diagnosis and requires further confirmatory diagnosis using molecular biology methods such as polymerase chain reaction (PCR), fluorescent *in-situ* hybridization (FISH) and sequencing. The evolution of molecular biology has made PCR-based diagnostic the most valuable method in identifying disease-causing pathogens to prevent and control diseases in bivalve culture. The development of PCR not only permits fast and sensitive screening of pathogens, but it also allows early detection of low infections (Pena, 2002). Apart from PCR, transmission electron microscope (TEM) is also highly recommended to identify and determine the structure of the viruses (Bai et al., 2016).

Marine bivalves are a reservoir of unique bioactive molecules for the development of alternative antimicrobials with AMPs being regarded as encouraging candidates for new and more environmentally friendly antimicrobials. Naturally-derived AMPs are suitable templates for the development of alternatives to antibiotics (Galdiero et al., 2015), and bivalve AMPs are shown to be structurally different from their analogue peptides obtained from the terrestrial habitat (Cheong et al., 2015). One of the most anticipated goals is the use of AMPs with better characteristics, like stronger antibacterial activity, lower cytotoxicity and ease of production for obtaining a marketable drug (Zanella et al., 2017).



Figure 5.3 A healthy oyster (left) versus an oyster infected with *Haplosporidium nelsoni* (right) (taken from https://barnegatshellfish.org/oyster_disease.htm)

5.5 Disease Management in Seaweed Culture

For centuries, seaweeds are widely consumed in Asia, especially in China (Yang et al., 2017), Japan and Korea (McHugh, 2003). Asian countries are the major consumers of seaweed products but these products are gaining popularity and are being exported to Europe, North America, and Africa (Chen & Xu, 2005). The popularity and beneficial properties of the seaweeds have triggered the massive expansion of the commercial production of seaweeds. Various species of seaweeds have been studied extensively to understand the potential use of seaweeds as dietary supplements, animal feed, chemicals

and biofuels (Tiwari & Troy, 2015). The seaweed aquaculture production is mainly focused within Asia, where it has dominated more than 80% of global production (Kim et al., 2017). Despite the tremendous development of the seaweed aquaculture technologies, this industry still struggles with disease outbreaks, causing the decline in seaweed yields.

Due to the economic importance of *Pyropia*, the best studied seaweed diseases are oomycete pathogens, *Pythium porphyrae* and *Olpidiopsis porphyrae*. Red rot disease caused by *P. porphyrae* was first recorded in Japan (Arasaki, 1947); however, it was only identified as the causative agent after three decades and recognised as the reason of the *Pyropia* crop yield reductions and quality (Takahashi et al., 1977). Initially, *Olpidiopsis* disease was assumed to be caused by a chytrid pathogen, but *Olpidiopsis* spp. was later identified as an obligate endoparasitic oomycete. *Olpidiopsis* disease seems to be endemic as it was only recorded in Japan (Arasaki, 1960; Sekimoto et al., 2008), Korea and China (Kim et al. 2014; Kwak et al., 2017). In Korea, the outbreak of oomycete pathogens is estimated to reduce up to 20% of *Pyropia* yields (Kim et al., 2014), with most of this loss attributed to *O. pyropiae* (Klochkova, et al., 2012) and green-spot disease, caused by the virus PyroV1 (Kim et al., 2016). Despite research efforts on the interactions between oomycete pathogens and their respective *Pyropia* hosts (Klochkova et al., 2012), these studies have offered little insight on biosecurity strategies that can be used in controlling and preventing disease outbreak in culture environments. Recent studies have shown that the red rot disease is not only caused by *P. poryphyrae* but also by another oomycete, *P. chondricola* (Qiu et al., 2019) and the fungus *Alternaria* sp. (Mo et al., 2016). Similarly, *O. pyropia* (Klochkova et al., 2016) has also been found to be the causative agent of *Olpidiopsis* disease, suggesting that the relationship between symptoms and agents may be more convoluted than known.

Ice-ice disease which affects *Kappaphycus* and *Euचेuma* spp., is characterized by thallus whitening in response to environmental stress such as changes in water temperature, salinity and pH, along with the action of opportunistic pathogenic bacteria (Ask & Azanza, 2002). In the past few decades, several bacterial interactions were found to be associated with ice-ice: *Cytophaga-Flavobacterium* complex (Largo et al., 1995) and *Alteromonas - Pseudoalteromonas* (Syafitri et al., 2017). Marine fungi such as *Aspergillus ochraceus*, *A. terreus* and *Phoma* sp. can induce ice-ice symptoms in *K. alvarezii* and *K. striatum* (Solis et al., 2010). Although the development of ice-ice symptoms results from stress to the host from abiotic conditions (Hurtado & Critchley, 2006; Vairappan et al., 2008) together with the action of opportunistic bacteria, the exact mechanisms that instigate the onset and progression of ice-ice symptoms remain unknown to date.

Apart from microorganisms, another significant pest of commercially cultivated seaweeds are epiphytic algae (Ward et al., 2019). Numerous studies have proven that the epiphytic filamentous algae (EFA) are responsible for a reduction in production biomass and carrageenan quality of cultivated *Kappaphycus* and *Euचेuma* spp. in Southeast Asia (Critchley et al., 2004; Hurtado & Critchley, 2006). EFA are found to occur in all major euचेumatoid-producing countries (Pang et al., 2015; Tsiresy et al., 2016). The declines in *K. alvarezii* production in Malaysia are caused by outbreaks of *Polysiphonia* (Hurtado & Critchley, 2006) and die-offs of the same crop species have been attributed to the presence

of *Melanothamnus savatieri* in Chinese farms since 2009 (Pang et al., 2011). The fluctuations of abiotic stress such as the seawater temperature, salinity, and current, and intensity of light and nutrient availability not only affect the host strength, but it could trigger the epiphyte infestations (Mtolera et al., 1996; Vairappan, 2006; Pang et al., 2011). (Pang et al., 2011) (Figure 5.4). Epiphytic algae infection causes damage to the cortex of *Kappaphycus* that leads to secondary bacterial infection (Vairappan et al., 2008).

Seaweed diseases have resulted in huge losses as exemplified in Indonesia, the world's largest seaweed producer (Kambey et al., 2020). Over the years, various treatment and mitigation strategies have been introduced to control the impact and spread of disease in the seaweed aquacultures. Acid wash is a commonly used treatment to control all diseases in the seaweed cultivation, particularly *Pyropia* (Kim et al., 2014). Studies have shown that immersion of *Pyropia* blades in acid solutions for more than 5 minutes could prevent the spread of red rot disease (Sakaguchi et al., 2001). This acid wash treatment has found to be partially effective for the red rot disease, cyanobacteria and diatom felt, but it is ineffective for the two most severe diseases of *Pyropia* crops: green-spot disease and *Olpidiopsis* disease (Kim et al., 2014). However, some studies have shown promising results with the use of bioactive metabolites produced by fungal endophytes in inhibiting the infection of *P. yezoensis* by *O. pyropiae* and *P. porphyrae* (Prado et al., 2017; Qiu et al., 2019). Additionally, the change of cultivation conditions with modifications in salinity and light exposure are effective measures in reducing the severity of diseases caused by exposure of *Saccharina japonica* to abiotic conditions (Wang et al., 2014).

Until today, the most effective technique in reducing the impact of pest epiphytes on seaweed production is the constant monitoring of the cultivated populations. This technique allows the farmer to remove those EFA infested crops immediately and replace them with healthy propagules from a different location (Ask & Azanza, 2002). However, there is still no effective management protocols for ice-ice disease to date. Before establishing a cost-effective management protocol, more studies are necessary to understand the underlying mechanisms behind the inset and the progression of ice-ice disease in *Kappaphycus* and *Euclima*.



Figure 5.4 A red algal epiphytes (left arrow) *Polysiphonia* and diatoms (right dotted arrow) on *Kappaphycus* thalli (adapted from Largo, 2002)

5.6 Conclusion

To conclude, it is important to remember that an alliance among three factors is necessary for a disease outbreak to happen. This alliance is the “epidemiological triangle” and consists of a host, a pathogen and the environment. It can be concluded that extreme temperatures, varying levels of salinity, human handling, high stocking densities in rearing systems and co-infection with other pathogens may reduce host immunity as well as increase the pathogen’s growth rate and infectivity, resulting in higher susceptibility to a disease. Based on knowledge and research on how a disease can spread whilst acknowledging the role of the environment, improving surveillance on the environment of rearing systems is very important. As knowledge of aquaculture diseases increase, it is important to switch the mindset of one-disease-one-pathogen paradigm to the “pathobiome” concept (Bass et al., 2019). “Pathobiome” systems consist of multiple host-associated organisms associated with decreased host health because of interactions between members of that set, the host, and environmental factors. The paradigm of disease management must shift to management of multiple organisms as well as the use of integrated measures for a healthy and sustainable ecosystem within the aquaculture ecosystem.

References

- Adams, A. (2016) *Fish Vaccines*, Springer Science+Business Media, Basel, Switzerland, 1st edition.
- Admodisastro, V.A., Ransangan, J., Ilias, N. and Tan, S.H. (2021). Oyster farming potential in Sabah, Malaysia. *International Journal of Aquatics Research*, 2, 17-22.
- Aly, S.M. and Albutti, A., (2014). Antimicrobials Use in Aquaculture and their Public Health Impact,” *J. Aquac. Res. Dev.*, vol. 5, no. 4.
- Arasaki, S. (1947). Studies on the wasting disease of the cultured lavers (*Porphyra tenera*). *Bull. Japan. Soc. Sci. Fish*, 13, 74–90
- Arasaki, S. (1960). A chytridean parasite on the *Porphyra*. *Nippon Suisan Gakkashi*, 26, 543–548.
- Ask, E. I., & Azanza, R. V. (2002). Advances in cultivation technology of commercial euclidean species: a review with suggestions for future research. *Aquaculture*, 26, 257–277.

- Assefa, A. and Abunna, F. (2018). Maintenance of Fish Health in Aquaculture: Review of Epidemiological Approaches for Prevention and Control of Infectious Disease of Fish. *Veterinary Medicine International*, Volume 2018, Article ID 5432497, 10 pages <https://doi.org/10.1155/2018/5432497>
- Bachère, E. (2003). Anti-infectious immune effectors in marine invertebrates: potential tools for disease control in larviculture. *Aquaculture* 227, 427–438.
- Bachere, E., Destoumieux, D. and Bulet, P. (2000). Penaeidins, antimicrobial peptides of shrimp: A comparison with other effectors of innate immunity. *Aquaculture*, 191, 71–88.
- Bai, C., Gao, W., Wang, C., Yu, T., Zhang, T., Qiu, Z., Wang, Q. and Huang, J. (2016) Identification and characterization of ostreid herpesvirus 1 associated with massive mortalities of *Scapharca broughtonii* broodstocks in China. *Dis. Aquat. Organ.* 2016, 118, 65–75.
- Bai, C., Wang, C., Xia, J., Sun, H., Zhang, S. and Huang, J. (2015) Emerging and endemic types of Ostreid herpesvirus 1 were detected in bivalves in China. *J. Invertebr. Pathol.*, 124, 98–106.
- Bass, D., Stentiford, G. D., Wang, H.-C., Koskella, B., & Tyler, C. R. (2019). The pathobiome: new insights in animal and plant health. *Trends Ecol. Evol.*, 34, 996-1008.
- Bene, C., Barang, M., Subasingh, R., Pinstруп-Andersen, P., Merino G, Hemr G.I. and Williams, M. (2015). Feeding 9 billion by 2050: Putting fish back on the menu. *Food Sec.*
- Bentzon-Tilia, M., Sonnenschein, E.C. and Gram, L. (2016). Monitoring and managing microbes in aquaculture – Towards a sustainable industry. *Microb. Biotechnol.*, 9,576–584.
- Bidault, A., Richard, G.G., le Bris, C. and Paillard, C. (2015) Development of a *Taqman* real-time PCR assay for rapid detection and quantification of *Vibrio tapetis* in extrapallial fluids of clams. *PeerJ*, 3, e1484.
- Biel, F.M., Allen, F.A. and Hase, C.C. (2014) Autolysis in *Vibrio tubiashii* and *Vibrio coralliilyticus*. *Can. J. Microbiol.*, 60, 57–63.
- Burreson, E.M. and Ragone Calvo, L.M. (1996). Epizootiology of *Perkinsus marinus* disease of oysters in Chesapeake Bay, with emphasis on data since 1985. *J. Shellfish Res.*, 15, 17–34.
- Carbone, D. and Faggio, C. (2016). Importance of prebiotics in aquaculture as immunostimulants. Effects on immune system of *Sparus aurata* and *Dicentrarchus labrax*. *Fish Shellfish Immunol.*, 54, 172–178.
- Chang, Y.-H., Kumar, R., Ng, T. H., & Wang, H.-C. (2017). What vaccination studies tell us about immunological memory within the innate immune system of cultured shrimp and crayfish. *Dev. Comp. Immunol.*, 80, 53–56.
- Chen, J., & Xu, P. (2005). Cultured aquatic species information programme, *Porphyra* spp. In FAO Fisheries and Aquaculture Department [online]. Rome
- Cheung, R.C.; Ng, T.B.; Wong, J.H. (2015) Marine peptides: Bioactivities and applications. *Mar. Drugs*, 13, 4006–4043
- Da Silva, P.M., Renaut, T., Fuentes, J., Villalba, A. (2008) Herpesvirus infection in European flat oysters *Ostrea edulis* obtained from brood stocks of various geographic origins and grown in Galicia (NW Spain). *Dis. Aquat. Organ.*, 78, 181–188
- Defoirdt, T., Sorgeloos, P. and Bossier, P. (2011). Alternatives to antibiotics for the control of bacterial disease in aquaculture. *Curr. Opin. Microbiol.*, 14, 251–258.
- Destoumieux-Garzón, D., Rosa, R.D., Schmitt, P., Barreto, C., Vidal-Dupiol, J., Mitta, G., Gueguen, Y., & Bachère, E. (2016). Antimicrobial peptides in marine invertebrate health and disease. *Philos. Trans. R. Soc. Lond., B, Biol. Sci.*, 371, 20150300.
- Dhar, A., Cowley, J., Hasson, K. and Walker, P. (2004). Genomic organization, biology, and diagnosis of Taura syndrome virus and yellowhead virus of penaeid shrimp. *Adv. Virus Res.*, 63, 353–421.
- FAO. Aquaculture Development Beyond (2000). The Bangkok Declaration and Strategy. Conference on Aquaculture in the Third Millennium, 20±25 February 2000. Bangkok. Thailand. NACA. Bangkok and FAO. Rome; 27p
- Farley, C.A., Banfield, W.G., Kasnic, G., Jr. and Foster, W.S. (1972) Oyster herpes-type virus. *Science*, 178, 759–760
- Fathi, S., Harun, A. N., Rambat, S. and Tukiran, N.A. (2018). Current Issues in Aquaculture: Lessons from Malaysia. *Adv. Sci. Lett.*, 24, 503-505.
- Fernandez Robledo, J.A., Vasta, G.R. and Record, N.R. (2014) Protozoan parasites of bivalve molluscs: Literature follows culture. *PLoS ONE*, 9, e100872
- Flegel, T.W. (2006). The special danger of viral pathogens in shrimp translocated for aquaculture. *Sci.*, 32, 215–231.
- Flegel, T.W. and Fegan, D. F. (2002). Strategies for preventing the spread of fish and shellfish diseases. *Fisheries Science*, 68, 776–788.
- Food and Agriculture Organization of the UN (FAO) (2012). The State of the World Fisheries and Aquaculture. 2012. FAO, Rome, Italy.
- Ford, S.E. (2011) Dermo Disease of Oysters Caused by *Perkinsus marinus*; Ford, S.E., Ed.; ICES Identification Leaflets for Diseases and Parasites of Fish and Shellfish; ICES: Copenhagen, Denmark.
- Francavilla, M., Franchi, M., Monteleone, M., & Caroppo, C. (2013). The red seaweed *Gracilaria gracilis* as a multi products source. *Mar. Drugs*, 11, 754–3776.
- Galdiero, S.; Falanga, A.; Berisio, R.; Grieco, P.; Morelli, G.; Galdiero, M. (2015) Antimicrobial peptides as an opportunity against bacterial diseases. *Curr. Med. Chem.*, 22, 1665–1677
- Guo, X. and Ford, S.E. (2016) Infectious diseases of marine molluscs and host responses as revealed by genomic tools. *Philos. Trans. R. Soc. Lond. B Biol. Sci.*, 371
- Huang, S., Wang, L., Liu, L., Hou, Y. and Li, L. (2015) Nanotechnology in agriculture, livestock, and aquaculture in China. A review. *Agronomy for Sustainable Development*, 35, 369–400.
- Hurtado, A. Q., & Critchley, A. T. (2006). Occurrence of Polysiphonia epiphytes in Kappaphycus farms at Calaguas Is., Camarines Norte, Philippines. *J. Appl. Phycol.*, 18, 301–306
- Itsathitphaisarn, O., Thitamadee, S., Weerachatanukul, W., & Sritunyalucksana, K. (2017). Potential of RNAi applications to control viral diseases of farmed shrimp. *Journal of Invertebrate Pathology*, 147, 76–85.
- Jennings, S. et al. (2016). Aquatic food security: trends, challenges and solutions for a single nation embedded in a dynamic global web of producers, processors and markets. *Fish Fish.*, 17, 893-938.

- Kambey, C.S.B., Campbell, I., Sondak, C.F.A., Nor, A.R.M., Lim, P.E. and Cottier-Cook, E.J. (2020). An analysis of the current status and future of biosecurity frameworks for the Indonesian seaweed industry. *J. Appl. Phycol.*, 32, 2147–2160
- Kim, G. H., Moon, K.-H., Kim, J.-Y., Shim, J., and Klochkova, T. A. (2014). A reevaluation of algal diseases in Korean *Pyropia* (*Porphyra*) sea farms and their economic impact. *Algae*, 29, 249–265.
- Kim, J. K., Yarish, C., Hwang, E. K., Park, M., and Kim, Y. (2017). Seaweed aquaculture: cultivation technologies, challenges and its ecosystem services. *Algae*, 32, 1–13.
- Klochkova, T. A., Shim, J. B., Hwang, M. S., and Kim, G. H. (2012). Host-parasite interactions and host species susceptibility of the marine oomycete parasite, *Olpidiopsis* sp., from Korea that infects red algae. *J. Appl. Phycol.*, 24, 135–144.
- Klochkova, T. A., Shin, Y. J., Moon, K.-H., Motomura, T., and Kim, G. H. (2016). New species of unicellular obligate parasite, *Olpidiopsis pyropiae* sp. nov., that plagues *Pyropia* sea farms in Korea. *J. Appl. Phycol.*, 28, 73–83.
- Kueh, C.S. & Chan, K.Y. (1985) Bacteria in bivalve shellfish with special reference to the oyster. *J. Appl. Bacteriol.*, 59, 41–47.
- Kumar, V., Roy, S., Meena, D.K. and Sarkar, U.K. (2016). Application of probiotics in shrimp aquaculture: importance, mechanisms of action, and methods of administration. *Rev. Fish. Sci. Aquac.*, 24 , 342–368.
- Labreuche, Y., and Warr, G. W. (2013). Insights into the antiviral functions of the RNAi machinery in penaeid shrimp. *Fish Shellfish Immunol.*, 34, 1002–1010.
- Lafferty, K.D., Harvell, C.D., Conrad, J.M., Friedman, C.S., Kent, M.L., Kuris, A.M. et al. (2015) Infectious diseases affect marine fisheries and aquaculture economics. *Ann. Rev. Mar. Sci.*, 7, 471–496.
- Largo, D. B., Fukami, K., & Nishijima, T. (1995). Occasional pathogenic bacteria promoting ice-ice disease in the carrageenan producing red algae *Kappaphycus alvarezii* and *Euचेuma denticulatum* (Solieriaceae, Gigartinales, Rhodophyta). *J. Appl. Phycol.*, 7, 545–554.
- Lee C-T, et al. (2015). The opportunistic marine pathogen *Vibrio parahaemolyticus* becomes virulent by acquiring a plasmid that expresses a deadly toxin. *Proc. Natl Acad. Sci. USA* 112, E5445.
- Leibovitz, L.; Elston, R.; Lipovsky, V.P. and Donaldson, J. (1978). A new disease of larval Pacific oysters (*Crassostrea gigas*). *J. World Aquacult. Soc.*, 9, 603–615.
- Leung, T.L.F. and Bates, A.E. (2013). More rapid and severe disease outbreaks for aquaculture at the tropics: Implications for food security. *J. Appl. Ecol.*, 50 , 215–222.
- Lightner, D. (1996). Epizootiology, distribution and the impact on international trade of two penaeid shrimp viruses in the Americas. *Rev. Sci. Tech.* 15, 579–601.
- Lightner, D.V. (1999). The penaeid shrimp viruses TSV, IHNV, WSSV, and YHV. *J. Appl. Aquac.*, 9, 27-52.
- Lightner, D. and Redman, R. (1981). A baculovirus-caused disease of penaeid shrimp, *Penaeus monodon*. *J. Invert. Pathol.*, 38, 299–302.
- Lim, P.-E., Yang, L.-E., Maggs, C. A., and Brodie, J. (2017). Advancing the taxonomy of economically important red seaweeds (Rhodophyta). *Eur. J. Phycol.*, 52, 438–451.
- Lo, C. et al. (1996). White spot syndrome baculovirus (WSBV) detected in cultured and captured shrimp, crabs and other arthropods. *Dis. Aquat. Organ.*, 27, 215–225.
- Lynch, S.A., Carlsson, J., Reilly, A.O., Cotter, E., Culloty, S.C. (2012) A previously undescribed Ostreid Herpes Virus 1 (OsHV-1) genotype detected in the Pacific oyster, *Crassostrea gigas*, in Ireland. *Parasitology*, 139, 1526–1532.
- McGladdery, S.E. (2011) Shellfish Diseases (Viral, Bacterial and Fungal) in Fish Diseases and Disorders Vol. 3: 748 Viral, Bacterial and Fungal Infections, 2nd Edition. P.T.K. Woo and D.W. Bruno Eds., pp748- 854. Cabi International,
- McHugh, D. J. (2003). A guide to the seaweed industry (FAO Fisheries Technical Paper 441). Rome, Italy: Food and Agriculture Organisation of the United Nations
- Miriam, R., Tapissier, N., Pierre, S. and Saulnier, D. (2017). Use of Medicinal Plants in Aquaculture, in Diagnosis and Control of Diseases of Fish and Shellfish, B. Austin and A. Newaj-Fyzul, Eds., pp. 223–261, JohnWiley & Sons Ltd., 1st edition.
- Mo, Z., Li, S., Kong, F., Tang, X., and Mao, Y. (2016). Characterisation of a novel fungal disease that infects the gametophyte of *Pyropia yezoensis* (Bangiales, Rhodophyta). *J. Appl. Phycol.*, 28, 395–404.
- Mortensen, S., Strand, A., Bodvin, T., Alfjorden, A., Skar, C.K., Jelmert, A., Aspan, A., Saelemyr, L., Naustvoll, L.J. and Alnbretsen, J. (2016) Summer mortalities and detection of ostreid herpesvirus microvariant in Pacific oyster *Crassostrea gigas* in Sweden and Norway. *Dis. Aquat. Organ.*, 117, 171–176.
- Moss, S. M., Moss, D. R., Arce, S. M., Lightner, D. V., and Lotz, J. M. (2012). The role of selective breeding and biosecurity in the prevention of disease in penaeid shrimp aquaculture. *J. Invertebr. Pathol.*, 110, 247–250.
- Munro, L.A. and Gregory, A. (2009) Application of network analysis to farmed salmonid movement data from Scotland. *J. Fish Dis.*, 32, 641–644.
- Murray, A.G. (2013) Epidemiology of the spread of viral diseases under aquaculture. *Current Opinion in Virology*, 3: 74–78.
- Naylor, R.L., Hardy, R.W., Buschmann, A.H., Bush, S.R., Cao, L., Klinger, D.H., Little, D.C., Lubchenco, J., Shumway, S.E. and Troell, M. (2021). A 20-year retrospective review of global aquaculture. *Nature*, 591, 551-563.
- Pang, T., Liu, J., Liu, J., and Li, J. (2015). Observations on pests and diseases affection a euचेumatoid farm in China. *J. Appl. Phycol.*, 27, 1975–1984.
- Pang, T., Liu, J., Liu, Q., and Lin, W. (2011). Changes of photosynthetic behaviours in *Kappaphycus alvarezii* infected by epiphytes. *Evidence-based Complementary and Alternative Medicine*, 2011, 658906. <https://doi.org/10.1155/2011/658906>
- Park, K.I., Yang, H.S., Kang, H.S., Cho, M., Park, K.J. and Choi, K.S. (2010) Isolation and identification of *Perkinsus olseni* from feces and marine sediment using immunological and molecular techniques. *J. Invertebr. Pathol.*, 105, 261–269.
- Paynter, K.T., Politano, V., Lane, H.A., Allen, S.M. & Meritt, D. (2010) Growth rates and prevalence of *Perkinsus marinus* in restored oyster populations in Maryland. *J. Shellfish Res.*, 29, 309–317.
- Pernet, F., Lupo, C., Bacher, C. & Whittington, R.J. (2016). Infectious diseases in oyster aquaculture require a new integrated approach. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* 371.
- Poulos, B., Tang, K., Pantoja, C., Bonami, J. and Lightner, D. (2006). Purification and characterization of infectious myonecrosis virus of penaeid shrimp. *J. Gen. Virol.*, 87, 987–996.

- Prado, S., Dubert, J., da Costa, F., Martinez-Patino, D. and Barja, J.L. (2014). Vibrios in hatchery cultures of the razor clam, *Solen marginatus* (Pulteney). *J. Fish Dis.*, 37, 209–217.
- Prado, S., Vallet, M., Gachon, C. M. M., Strittmatter, M., & Kim, G. H. (2017). Prevention or treatment in algae of diseases induced by protistan pathogens. Pub. No.: WO/2017/125775 [patent].
- Qiu, L., Mao, Y., Tang, L., Tang, X., and Mo, Z. (2019). Characterisation of *Pythium chondricola* associated with red rot disease *Pyropia yezoensis* (Ueda) (Bangiales, Rhodophyta) from Lianyungang, China. *J. Oceanol. Limnol.*, 37, 1102–1112.
- Renault, T. & Novoa, B. (2004). Viruses infecting bivalve molluscs. *Aquat. Living Resour.*, 17, 397–409.
- Robalino, J. et al. (2005). Double-stranded RNA induces sequence-specific antiviral silencing in addition to nonspecific immunity in a marine shrimp: Convergence of RNA interference and innate immunity in the invertebrate antiviral response? *J. Virol.*, 79, 13561–11371.
- Rogers, C. and Basurco, B. (2009). Antimicrobial agents in aquaculture: Practice, needs and issues,” *CIHEAM*, 59, pp. 41–59.
- Rogers, L.A., Bateman, A.W., Connors, B.M., Frazer, L.N., Godwin, S.C., Krkosšek, M., Lewis, M.A., Peacock, S.J., Rees, E.E., Revie, C.W. and Schlaegel, U.E. (2016). Lessons from sea louse and salmon epidemiology. *Phil. Trans. R. Soc. B*; 371, 20150203.
- Romero, J., Feijóo, C.G. and Navarrete, P. (2012). Antibiotics in aquaculture Use, abuse and alternatives. *Health and Environment in Aquaculture*, pp. 160–198.
- Sánchez-Paz, A. (2010). White spot syndrome virus: an overview on an emergent concern. *Vet. Res.* 41, 43 (10.1051/vetres/2010015)
- Segarra, A., Pepin, J.F., Arzul, I., Morga, B., Faury, N. and Renault, T. (2010) Detection and description of a particular Ostreid herpesvirus 1 genotype associated with massive mortality outbreaks of Pacific oysters, *Crassostrea gigas*, in France in 2008. *Virus Res.*, 153, 92–99
- Shekarm, M., Pradeep, B. & Karunasagar, I. (2012). White spot syndrome virus: Genotypes, epidemiology and evolutionary studies. *Indian J. Virology*, 23, 175-183.
- Shen, Y., Ma, K. and Yue, G.H. (2020) Status, challenges and trends of aquaculture in Singapore. *Aquaculture*, <https://doi.org/10.1016/j.aquaculture.2020.736210>
- Smolowitz, R. (2013). A review of current state of knowledge concerning *Perkinsus marinus* effects on *Crassostrea virginica* (the eastern oyster). *Vet. Pathol.*, 50, 404–411.
- Stentiford, G.D., Sritunyalucksana, K., Flegel, T.W., Williams, B.A.P., Withyachumnarnkul, B., Itsathitphaisarn, O. et al. (2017). New Paradigms to Help Solve the Global Aquaculture Disease Crisis. *PLoS Pathog* 13(2): e1006160.
- Sugumar, G., Nakai, T., Hirata, Y., Matsubara, D. and Muroga, K. (1998). *Vibrio splendidus* biovar II as the causative agent of bacillary necrosis of Japanese oyster *Crassostrea gigas* larvae. *Dis. Aquat. Organ.*, 33, 111–118.
- Stentiford, G.D. et al. (2012). Disease will limit future food supply from the global crustacean fishery and aquaculture sectors. *J. Invertebr Pathol.*, 110, 141-157.
- Takahashi, M., Ichitani, T., & Sasaki, M. (1977). *Pythium porphyrae* Takahasi et Sasaki, sp. nov. causing red rot of marine red algae *Porphyra* spp. *Trans. Mycol. Soc. Japan.*, 18, 279–285.
- Tan, S.H (2021). A BRIGHT FUTURE AWAITS OYSTER FARMING IN PENANG, Penang Monthly. Retrieved from <https://news.usm.my/index.php/rencana-features/6973-a-bright-future-awaits-oyster-farming-in-penang>. Accessed on 24 May 2022.
- Tan, S.H. (2018). Government Efforts to Promote Oyster Farming in Malaysia. *World Aquaculture*, 39-41 Department of Fisheries Malaysia, 2018. *Annual fisheries statistic 2018*. Retrieved 14/8/2020 from <http://www.dof.gov.my/en/fisherystatistics>
- Tassanakajon, A., Amparyup, P. and Premruethai, K.S. (2010). Cationic antimicrobial peptides in penaeid shrimp. *Mar. Biotechnol.*, 12, 487-505.
- Thitamadee, S., Prachumwat, A., Srisala, J., Sritunyalucksana, K., Flegel, T.W. and Itsathitphaisarn, O. (2016). Review of current disease threats for cultivated penaeid shrimp in Asia. *Aquaculture*, 452, 69–87.
- Tiwari, B. K., and Troy, D. J. (Eds.). (2015). *Seaweed sustainability: Food and non-food applications*. Cambridge, Massachusetts: Academic Press, 472 pp
- Tsiresy, G., Preux, J., Latriva, T., Dubois, P., Lepoint, G., and Eeckhaut, I. (2016). Phenology of farmed seaweed *Kappaphycus alvarezii* infestation by the parasitic epiphyte *Polysiphonia* sp. in Madagascar. *J. Appl. Phycol.*, 28, 2903–2914.
- Tubiash, H.S., Chanley, P.E. and Leifson, E. (1965). Bacillary necrosis, a disease of larval and juvenile bivalve mollusks. I. Etiology and epizootiology. *J. Bacteriol.*, 90, 1036-1044.
- Vairappan, C. S., Chung, C. S., Hurtado, A. Q., Msuya, F. E., Bleicher Lhonneur, G., & Critchley, A. (2008). Distribution and symptoms of epiphyte infection in major carrageenophyte-producing farms. *J. Appl. Phycol.*, 20, 477–483.
- Valente, C.S. & Wan, A.H.L. (2021). *Vibrio* and major commercially important vibriosis diseases in decapod crustaceans. *Journal of Invertebrate Pathology*, 181.
- Valerio, E., Chaves, S. and Tenreiro, R. (2010) Diversity and impact of prokaryotic toxins on aquatic environments: A review. *Toxins* (Basel), 2, 2359–2410
- Walker, P.J. and Mohan, C.V. (2009). Viral disease emergence in shrimp aquaculture: origins, impact and the effectiveness of health management strategies. *Rev. Aquac.*, 1, 125-154.
- Ward, G.M., Faisan, J.P., Cottier-Cook, E.J., Gachon, C., Hurtado, A.Q., Lim, P.E., Matoju, I., Msuya, F.E., Bass, D. and Brodie, J. (2019). A review of reported seaweed diseases and pests in aquaculture in Asia. *J. World Aquacult. Soc.*, 51, 815–828.
- Yang, L.-E., Lu, Q. Q., & Brodie, J. (2017). A review of the bladed Bangiales (Rhodophyta) in China: history, culture and taxonomy. *Eur. J. Phycol.*, 52, 1–13.
- Yodmuang, S., Tirasophon, W., Roshorm, Y., Chinnirunvong, W., and Panyim, S. (2006). YHV-protease dsRNA inhibits YHV replication in *Penaeus monodon* and prevents mortality. *Biochem. Biophys. Res. Commun.*, 341, 351–356.
- Zannella, C., Mosca, F., Mariani, F., Franci, G., Folliero, V., Galdiero, M., Tiscar, P.G. and Galdiero, M. (2017) Microbial Diseases of Bivalve Mollusks: Infections, Immunology and Antimicrobial Defense. *Mar. Drugs*, 15, 182.

Chapter 6

CURRENT UPDATE ON THE STRATEGIES USED IN COMBATING ACUTE HEPATOPANCREATIC NECROSIS DISEASE (AHPND)

Wen-Chian Tan¹, Hok-Chai Yam^{1,2}*

¹ Faculty of Applied Sciences, UCSI University, 56000 Cheras, Kuala Lumpur, Malaysia.

² Centre of Research for Advanced Aquaculture (CORAA), UCSI University, 56000, Cheras, Kuala Lumpur, Malaysia.

*Corresponding author: yamhc@ucsiuniversity.edu.my

ABSTRACT

Acute hepatopancreatic necrosis disease (AHPND) is a shrimp disease primarily caused by virulent *Vibrio* species. AHPND has had a devastating directed against the shrimp farming industry with an approximate plunge to 60 % production and has resulted in collective losses of more than USD 43 billion in Asia. This chapter provides an update on the practices being used to mitigate the impact of AHPND and the ongoing research findings that potentially contribute to new strategies.

KEYWORDS: Acute hepatopancreatic necrosis disease, early mortality syndrome, herbs in aquaculture, probiotic in aquaculture, white leg shrimp.

6.1 Introduction

Vibrio species, that cause acute hepatopancreatic necrosis disease (AHPND), have escalated since the South Asian country collapsed in late 2013. AHPND has a huge negative impact on the shrimp industry and grows rapidly, starting approximately eight days after stocking at postlarvae stage 8 (PL 8), with severe mortality (up to 100 %) occurring within 20 - 30 days (PL 20 - PL 30). As a result, shrimp

production in the AHPND-affected areas has temporarily dropped to about 60 % and led to collective losses of more than USD 43 billion across Asia.

Most treatments and control measures developed for AHPND have focused on *Vibrio parahaemolyticus* (VP). However, the presence of a plasmid of AHPND-causing pVA1 with the size of 63–70 kb, encoding toxins PirAVP and PirBVP in non-VP and even non-*Vibrio* species have raised concerns. In addition, the toxin gene seems to migrate horizontally between bacteria (Dong et al., 2017). Hence, the causative agent for AHPND is no longer limited to single species of bacteria. Also, VP cells can replicate in less than fifteen minutes, while other *Vibrio* species take more than an hour.

As the causative agent could be more than one bacteria, investigating management practices to mitigate specific VP is not practical and would put unnecessary economic stress on farmers. Hence, control measures based on the existence of PirABVP toxins in shrimp and aquaculture systems may be more appropriate for the eradication of AHPND in shrimp culture systems. Also, there is interest in investigating methods that can enhance the innate immune responses and enhance disease resistance in shrimp.

In aquaculture, antibiotics is the most common method of treating outbreaks caused by *Vibrio* species; however, this led to the rise of antibiotic resistance, as is the possible spread of drugs in the environment. Consequently, other methods, for instance, disinfection before shrimp post-larvae stocking, pond management, vaccines, supplementation of immunostimulants, prebiotics, probiotics or phages, have gained the interest of farmers in combating AHPND. Nevertheless, there are advantages and disadvantages that farmers need to consider before choosing a method. Therefore, this monograph presents the current strategies and potential methods for treating this disease from the researcher's perspective.

6.2 Current Strategies and Treatments

6.2.1 Antibiotics

VP, which causes early mortality syndrome in white leg shrimp, is a food-borne pathogen caused by bacteria that may cause human gastroenteritis. The bacteria can transmit to humans by consumption of contaminated seafood. Ingestion of VP-contaminated food can cause serious gastrointestinal illness (Tan et al., 2020). To overcome the scenario, antibiotics as feed additives are frequently used in prevention or treatment (Devi et al., 2009).

The use of antibiotics disrupts microbial maturation systems leading to ineffectiveness in treating diseases caused by *Vibrio* spp. (de Schryver et al., 2014). Anyway, this will lead to another serious threat: multidrug-resistance. In a study by Tan et al. (2020), VP prevalence and antibiotic resistance in blood clams, shrimp, jellyfish and squid were observed. Another research by Lai et al. (2015) concludes that both virulent and non-virulent VP strains are resistant to several antibiotics commonly used in

shrimp farming. Nevertheless, inappropriate use of antibiotics in combating AHPND can lead to the development of VP resistant strains. At the same time, *V. campbellii* (that emerged from China), was found to have several genes that encodes with antibiotic resistance (Dong et al., 2017), so traditional methods such as antibiotics were used for shrimp farming. In addition, Dong et al. (2017) revealed that the spread of risk genes in *Vibrio* spp. is likely due to horizontal interspecies gene transfer, allowing VPAHPND to spread between the animals and water, to accelerate plague epidemic disease. Research from (Devi et al., 2009) reveals that a higher total putative VP count and antibiotic resistance were found in pond sediment samples, especially during the pre-monsoon season. Hence, this phenomenon suggests that antibiotic use in shrimp farming should be more strictly regulated.

6.2.2 Disinfection Before Shrimp Postlarvae Stocking

Once AHPND appears in ponds or adjacent ponds, disinfection does not stop the epidemic, so more effective treatments are immediately needed to control AHPND in shrimp. Traditional methods for alleviating or curing VPAHPND strains, for example, alternate feeding or using antibiotics and disinfectants, have had limited benefit (Hong et al., 2016). Thorough bottom of the pond disinfection and water to eliminate possible VPAHPND may help reduce the epidemic outbreak of AHPND disease instead of controlling it, and pond management based on microbial control strategies and ecological theory may be essential (de Schryver et al., 2014). Pond management, that involve shrimp culture in microbes' ecosystem balance ecosystems and applying biological control method that in-line with these procedures, are critical in addressing these issues (de Schryver et al., 2014). Furthermore, their use in the food production sector is subject to rigorous research and general scrutiny due to the expansion of multidrug resistance (Cao et al., 2021). For example, VPAHPND from Mexico was reported to carry the *tetB* gene encoding a tetracycline resistance gene, which shows high levels of resistance to tetracyclines (≥ 5 g/mL) (Han et al., 2015).

6.2.3 Immune Priming and Quorum Sensing

Immune priming is a memory-like phenomenon described in the invertebrate taxa of animals. For example, using inactivated pathogens against shrimp disease has been reported to induce memory responses in shrimp (Fu et al., 2011). Shrimp-specific immune priming makes "vaccination" desirable and feasible (Fu et al., 2011). Furthermore, it is evolutionarily beneficial for organisms to develop better and faster secondary immune responses to pathogens, which are detrimental and likely to be re-exposed.

Next, invertebrate shrimp are thought incomplete with proper lymphocyte and upper humoral immunity response. Instead, they depend primarily on innate or nonspecific immunity. Immune priming is defined as the two-step vaccination technique in which a pathogen is administrated into the host and revaccinated. For example, upon introducing recombinant viral proteins from White spot syndrome virus (WSSV) into *Bacillus subtilis* spores, immune priming in PLs resulted in huge phagocytic, which is a key biological activity by which the host protects itself from infectious and non-infectious environmental particles and removes unwanted host cells to maintain tissue homeostasis against WSSV (Fu et al., 2011).

Immunity to vibriosis in shrimp vaccinated with pre-formalin-killed *Vibrio* species has a more prolonged postpartum survival (Nguyen et al., 2017). A better immunity to toxin pathogens originated from bacterial, observed in PV post-larvae or larger larvae. Resistance towards the pathogens of antigenic origin initially suggests that the protection elicited by immunization is only for a specific pathogen type and strain in the initial vaccine. Immune specificity is sensitive towards purely harmful bacteria that would harm the host.

On the other hand, antiviral therapy is a promising approach because it only disarms pathogens that affect virulence gene expression instead of killing them. For example, research from de Schryver et al. (2014) reported that element that inhibit quorum sensing and bacterial communication that modulates the expression of virulence factors reduce mortality in giant river shrimp larvae.

6.3 Disadvantages of Current Strategies: Antibiotic Resistance in Shrimp Bacterial Pathogens

The abused widespread use of antibiotics, insecticides and sterilize in aquaculture caused growth of drug-resistant bacteria and caused social concern (Kang et al., 2018) in aquaculture. Research from Kang et al. (2018) isolated and identified 59 VPs (toxR-positive) in shellfish fishing areas on the west coast of Korea between May 2016 and October 2016. The results showed that none of the isolates delimited the *tdh* and *trh* virulence genes. The multiple antibiotic resistance mean value of utmost isolates was 0.32, and the highest value was 0.69 in one isolate. Antibiotics are used in aquaculture as feed additives and baths to treatment and prevent the transmit of disease (Rico et al., 2013). Also, VP isolates in some aquatic products from Asian and European countries (e.g., southern China (Xie et al., 2020) have high resistance rates to antibiotics such as ampicillin, rifampicin, and streptomycin (Kang et al., 2018).

6.4 Research Progress and Potential Therapeutic or Control Agents

6.4.1 Probiotics

Probiotics are familiar as prevalent health products for humans, also used in farming. In aquaculture, beneficial bacterial have been broadly used to restrain the evolution of opportunistic pathogens and reduce the prevalence of viruses (Fu et al., 2011). Probiotic usage increased with the need for more environmentally friendly aquaculture practices addition, and probiotics can improve nutrients, enzymatic activity, immune responses, and water quality (Zhou et al., 2009). Using probiotics to inhibit VPAHPND infection or other bacteria in shrimp is practical and improves aquaculture water quality (Kumar et al., 2021). Healthy shrimp can produce digestive enzymes with a healthy gut, and this is often associated with the good expelling of pathogenic bacteria, also in production of elements beneficial for host health (Wang et al., 2020). However, most profit-making available probiotics for aquaculture come from surface-dwelling animals. These non-host are associated with more growing bacteria strains and are

relatively void because they cannot stay alive at the intense cell density of the host as well as overcome antagonism during the active growth phase (gut environment) of the shrimp.

6.4.1.1 *Bacillus* spp.

Lim et al. (2020) stated that shrimp intestine microbiota in the wild showed a positive antibacterial effect against VPAHPND. In addition, two isolates showed positive inhibition against VPAHPND, namely *Penicillium* isolate B2 and *Penaeus monodon* isolate BT. In detailed, the isolates B2 and BT are represented as *Bacillus* species based on 16S rDNA sequencing results. The abovementioned bacterial isolates can survive in bile environments, exhibit excellent attachment to the intestinal mucus of shrimp, and are antibiotic-sensitive, non-hemolytic, and non-pathogenic to the host.

Research from Kewcharoen & Srisapoom (2019) isolated probiotic candidates approach coastal areas, and *Bacillus* were tested for the level in mitigating the harmfulness of VPAHPND strains. In addition, *Bacillus subtilis* AQAHBS001, which broadly inhibits various VPAHPND strains, was evaluated on a laboratory scale for water solubility and feed applications of live probiotics. Also, Zhou et al. (2009) showed that the additive of probiotic *Bacillus coagulans* SC8168 taken from shrimp pond sediment in Zhejiang Province, China, could significantly improve PV shrimp's survival rate and partial digestive enzyme (protease, amylase and lipase) activity.

6.4.1.2 Mixed Probiotics

Research from Chumpol et al. (2017) suggested that applying purple non-sulfur bacteria (PNSB) in white shrimp culture would help. Specifically, *Rhodobacter sphaeroides* strains and *Aifella marina* STW181 are beneficial bacterial for shrimp. However, throughout the research, Chumpol et al. (2017) observed a strong negative correlation between *Vibrio* numbers and shrimp survival. Conversely, a positive correlation was found between PNSB counts and shrimp survival. Another study by Nimrat et al. (2012) examined the efficacy of mixed *Bacillus* probiotic forms and probiotic applications on growth, showing that bacterial counts intersect with water quality LV shrimp culture in separate trials at the larval postlarval stages. The result showed that the form of probiotics and the mode of administration of the probiotics is harmless on the growth and survival of PL shrimp larvae (Nimrat et al., 2012). In addition, *Bacillus* microcapsules and freeze-dried probiotics expressively improved the growth and survival of PL shrimp only. Concerning this, the efficacy of probiotics might depend on exposure time at different stages of individual development (Ziaei-Nejad et al., 2006). Hence, the effectiveness of probiotics will be lesser in the larval period (4 days) into PL stages (22 days).

6.4.2 Phage Therapy

Phage therapies are promising approaches to preventing and treating vibriosis in aquaculture. In addition, phage therapy is a potential management strategy for controlling infectious diseases in humans and animals. The study by Lomelí-Ortega & Martínez-Díaz, (2014) evaluated the efficacy of bacteriophages in preventing and controlling vibriosis in shrimp.

Briefly, VP-infected *Penaeus vannamei* (PV) larvae receive with different phage doses, and the variable used was named phage A3S and Vpms 1. Researchers concluded that one single amount of Vpms1 or A3S phage is sufficient to minimize infection and dying rate in shrimp larvae lead by VP. The outstanding result has overwritten the previous effect from Karunasagar et al., (2019.) and Vinod et al. (2006), where a repeated dose is needed to treat AHPND bacteria phage naturally and is relatively inexpensive, without severe or irreversible side effects reported to date. With this, phage therapy was proven to prevent death by *Vibrio* effectively. In addition, the action of bacterio-phages is very selective, where some can recognize and infect only specific strains, compared to the use of antibiotics. This property is critical to ensure the effectiveness of phages as controls or preventive treatments for the disease in animals (Pereira et al., 2021) because mixed infections are common in cultured organisms. Exciting research from Cao et al. (2021) suggested using multidrug-resistant VP strains as bacteriophages that work as antibacterial agents to control AHPND. Therefore, the study isolated VP phage (vB_VpaP_MGD2) from the clam (*Meretrix meretrix*), targeting multidrug-resistant VP and further characterized it to assess its potential for biocontrol (Cao et al., 2021).

Jun et al. (2016) showed that bacteriophage pVp-1 exhibited significant bacteriolytic activity against three representative VPAHPNDs, resulting in 100% mortality within 24 hours of infection. The mortality of shrimp treated with phage was lower than that of the positive control group. In addition, hepatopancreas of phage-treated shrimp showed a marked region of AHPND with pathological changes after prophylactic treatments. The timing of phage addition concerning pathogen development is critical in vibriosis, which can be addressed by more frequent phage administration.

6.4.3 Plant-Derived Natural Compounds

Herb extracts were studied as an alternative to sustainable aquaculture. The research from Harlina et al. (2022.) focused on the antibacterial potential of *Tribulus terrestris* and basil against *Vibrio harveyi*, *V. alginolyticus* and VP. Research suggested that the organic acids blend (OAB) can significantly increase yield and interspersed to pathogenic *Vibrio*. The action may be a feasible alternative to antibiotics used in the shrimp industry because it exhibits hepatopancreas' protective properties against *Vibrio* spp. (Romano et al., 2015). The OAB includes *Piper betle* and *Cymbopogon citratus* extracts incorporated in their normal diet and showed good latent for treating diseases and colds, minimizing AHPND. Research from Immanuel et al. (2004) suggested that certain Ayurvedic herbs compounds have potent effects on the survival of aquatic life and exert specific antibacterial properties. The same studies have shown castor oil, *Phyllanthus niruri*, *Leucus aspera*, *Manihotesculenta* and seaweeds (*Ulva lactuca* and *Sargassum wightii*) can improve protect shrimps from VP infection (Immanuel et al., 2004).

6.4.4 Vaccine

Further research is focused on developing a vaccine against VPAHPND. In general, shrimp will prefer to have passive immunization due to the lack of an adaptive immune system. Specifically, kinetic immunity can protect shrimp from *Vibrio* infection by oral administration of a specific egg yolk powder (IgY) that inhibits VP and *Vibrio harveyi* in vitro (Gao et al., 2016). Relevant research has done just that, demonstrating that specific IgY effectively inhibits the growth of VP, providing passive immunity for shrimp (Hu et al., 2019). Recently, VP declined by the knockout of the LpxDgene gene that caused bacterial virulence and provided substantial protection to shrimp against challenges (Tsai et al., 2019). This method works with attenuated virulent bacteria as live attenuated vaccines against VP. Toxins A and B, since VP can cause AHPND, monoclonal antibodies to toxins A and B that can be used to detect toxins using dot blot (Wangman et al., 2017). Several studies used an anti-Pir A-like toxin IgY diet and demonstrated passive immunity against AHPND (Nakamura et al., 2019). Supplementing anti-Pir A-IgY in feeds be an effective method to prevent AHPND infection. Vaccine development against AHPND is a significant concern as further research is needed to understand the underlying mechanisms and pathogenesis of the disease.

6.5 Conclusion

Shrimp aquaculture is one of the intensified food production sectors in the world. However, the acute hepatopancreatic necrosis disease (AHPND) outbreak has resulted in substantial economic losses to the farming industry since 2009. The causative agent is the virulent *Vibrio* strains, including *V. parahaemolyticus*, *V. punensis*, *V. harveyi*, *V. owensey*, *V. campbellian* and *Shewanella* spp. that contains pVA1 plasmids (63–70 kb) encoding the binary PirAVP and PirBVP toxins. AHPND-affected shrimps exhibited notable histopathological changes, including massive shedding of hepatopancreatic epithelial cells without any accompanying signs of pathogens, suggesting that the binary PirAVP and PirBVP toxins secreted by bacteria are involved in the induction of AHPND. Furthermore, recent studies have shown that AHPND-related strains have other specific virulence factors involved in the virulence and disease pathology of AHPND-causing bacteria. Further molecular, microbiological, and histopathological studies are demanded to elucidate the virulence factors of AHPND-causing bacteria. This would help to develop newer strategies for mitigating the disease since most of the methods are still immature to be translated into practical use.

REFERENCES

- Cao, Y., Zhang, Y., Lan, W., & Sun, X. (2021). Characterization of vB_VpaP_MGD2, a newly isolated bacteriophage with biocontrol potential against multidrug-resistant *Vibrio parahaemolyticus*. *Archives of Virology*, 166(2), 413–426. <https://doi.org/10.1007/s00705-020-04887-x>
- Chumpol, S., Kantachote, D., Nitoda, T., & Kanzaki, H. (2017). The roles of probiotic purple nonsulfur bacteria to control water quality and prevent acute hepatopancreatic necrosis disease (AHPND) for enhancement growth with higher survival in white shrimp (*Litopenaeus vannamei*) during cultivation. *Aquaculture*, 473, 327–336. <https://doi.org/10.1016/j.aquaculture.2017.02.033>
- de Schryver, P., Defoirdt, T., & Sorgeloos, P. (2014). Early Mortality Syndrome Outbreaks: A Microbial Management Issue in Shrimp Farming? *PLoS Pathogens*, 10(4). <https://doi.org/10.1371/journal.ppat.1003919>

- Devi, R., Surendran, P. K., & Chakraborty, K. (2009). Antibiotic resistance and plasmid profiling of *Vibrio parahaemolyticus* isolated from shrimp farms along the Southwest coast of India. *World Journal of Microbiology and Biotechnology*, 25(11), 2005–2012. <https://doi.org/10.1007/s11274-009-0101-8>
- Dong, X., Bi, D., Wang, H., Zou, P., Xie, G., Wan, X., Yang, Q., Zhu, Y., Chen, M., Guo, C., Liu, Z., Wang, W., & Huang, J. (2017). pirABvp-Bearing *Vibrio parahaemolyticus* and *Vibrio campbellii* pathogens isolated from the Same AHPND-affected pond possess highly similar pathogenic plasmids. *Frontiers in Microbiology*, 8(OCT). <https://doi.org/10.3389/fmicb.2017.01859>
- Fu, L. L., Wang, Y., Wu, Z. C., & Li, W. F. (2011). In vivo assessment for oral delivery of *Bacillus subtilis* harboring a viral protein (VP28) against white spot syndrome virus in *Litopenaeus vannamei*. *Aquaculture*, 322–323, 33–38. <https://doi.org/10.1016/j.aquaculture.2011.09.036>
- Gao, X., Zhang, X., Lin, L., Yao, D., Sun, J., Du, X., Li, X., & Zhang, Y. (2016). Passive immune-protection of *Litopenaeus vannamei* against *Vibrio harveyi* and *Vibrio parahaemolyticus* infections with Anti-*Vibrio* Egg Yolk (IgY)-encapsulated feed. *International Journal of Molecular Sciences*, 17(5). <https://doi.org/10.3390/ijms17050723>
- Han, J. E., Mohnney, L. L., Tang, K. F. J., Pantoja, C. R., & Lightner, D. v. (2015). Plasmid mediated tetracycline resistance of *Vibrio parahaemolyticus* associated with acute hepatopancreatic necrosis disease (AHPND) in shrimps. *Aquaculture Reports*, 2, 17–21. <https://doi.org/10.1016/j.aqrep.2015.04.003>
- Harlina, H., Hamdillah, A., Rosmiati, R., Kasnir, M., & Syahrul, S. (2022). Potency of the Piper betle and *Ocimum basilicum* as a natural antibacterial against the acute hepatopancreatic necrosis disease (AHPND). *Aquaculture, Aquarium, Conservation & Legislation*, 15(1), 34-43.
- Hong, X., Lu, L., & Xu, D. (2016). Progress in research on acute hepatopancreatic necrosis disease (AHPND). In *Aquaculture International* (Vol. 24, Issue 2, pp. 577–593). Springer International Publishing. <https://doi.org/10.1007/s10499-015-9948-x>
- Hu, B., Yang, X., Guo, E., Zhou, P., Xu, D., Qi, Z., & Deng, L. (2019). The preparation and antibacterial effect of egg yolk immunoglobulin (IgY) against the outer membrane proteins of *Vibrio parahaemolyticus*. *Journal of the Science of Food and Agriculture*, 99(5), 2565–2571. <https://doi.org/10.1002/jsfa.9470>
- Immanuel, G., Vincybai, V. C., Sivaram, V., Palavesam, A., & Marian, M. P. (2004). Effect of butanolic extracts from terrestrial herbs and seaweeds on the survival, growth and pathogen (*Vibrio parahaemolyticus*) load on shrimp *Penaeus indicus* juveniles. *Aquaculture*, 236(1–4), 53–65. <https://doi.org/10.1016/j.aquaculture.2003.11.033>
- Jun, J. W., Han, J. E., Tang, K. F. J., Lightner, D. v., Kim, J., Seo, S. W., & Park, S. C. (2016). Potential application of bacteriophage pVp-1: Agent combating *Vibrio parahaemolyticus* strains associated with acute hepatopancreatic necrosis disease (AHPND) in shrimp. *Aquaculture*, 457, 100–103. <https://doi.org/10.1016/j.aquaculture.2016.02.018>
- Kang, C. H., Shin, Y. J., Yu, H. S., Kim, S. K., & So, J. S. (2018). Antibiotic and heavy-metal resistance of *Vibrio parahaemolyticus* isolated from oysters in Korea. *Marine Pollution Bulletin*, 135, 69–74. <https://doi.org/10.1016/j.marpolbul.2018.07.007>
- Karunasagar, I., Vinod, M. G., Kennedy, B., Vijay, A., Deepanjali, A., Umesh, K. R., & Karunasagar, I. (n.d.). *Biocontrol of Bacterial Pathogens in Aquaculture with Emphasis on Phage Therapy*. <http://www.who.int/inf-fs/en/fact>
- Kewcharoen, W., & Srisapoome, P. (2019). Probiotic effects of *Bacillus* spp. from Pacific white shrimp (*Litopenaeus vannamei*) on water quality and shrimp growth, immune responses, and resistance to *Vibrio parahaemolyticus* (AHPND strains). *Fish and Shellfish Immunology*, 94, 175–189. <https://doi.org/10.1016/j.fsi.2019.09.013>
- Kumar, V., Roy, S., Behera, B. K., Bossier, P., & Das, B. K. (2021). Acute hepatopancreatic necrosis disease (Ahpnd): Virulence, pathogenesis and mitigation strategies in Shrimp aquaculture. In *Toxins* (Vol. 13, Issue 8). MDPI AG. <https://doi.org/10.3390/toxins13080524>
- Lai, H. C., Ng, T. H., Ando, M., Lee, C. te, Chen, I. T., Chuang, J. C., Mavichak, R., Chang, S. H., Yeh, M. de, Chiang, Y. A., Takeyama, H., Hamaguchi, H. o., Lo, C. F., Aoki, T., & Wang, H. C. (2015). Pathogenesis of acute hepatopancreatic necrosis disease (AHPND) in shrimp. *Fish and Shellfish Immunology*, 47(2), 1006–1014. <https://doi.org/10.1016/j.fsi.2015.11.008>
- Lim, S. Y., Loo, K. W., & Wong, W. L. (2020). Synergistic Antimicrobial Effect of a Seaweed-Probiotic Blend Against Acute Hepatopancreatic Necrosis Disease (AHPND)-Causing *Vibrio parahaemolyticus*. *Probiotics and Antimicrobial Proteins*, 12(3), 906–917. <https://doi.org/10.1007/s12602-019-09616-8>
- Lomeli-Ortega, C. O., & Martínez-Díaz, S. F. (2014). Phage therapy against *Vibrio parahaemolyticus* infection in the whiteleg shrimp (*Litopenaeus vannamei*) larvae. *Aquaculture*, 434, 208–211. <https://doi.org/10.1016/j.aquaculture.2014.08.018>
- Nakamura, A. (n.d.). 2 |.
- Nguyen, H. T., Thu Nguyen, T. T., Tsai, M. A., Ya-Zhen, E., Wang, P. C., & Chen, S. C. (2017). A formalin-inactivated vaccine provides good protection against *Vibrio harveyi* infection in orange-spotted grouper (*Epinephelus coioides*). *Fish and Shellfish Immunology*, 65, 118–126. <https://doi.org/10.1016/j.fsi.2017.04.008>
- Nimrat, S., Suksawat, S., Boonthai, T., & Vuthiphandchai, V. (2012). Potential *Bacillus* probiotics enhance bacterial numbers, water quality and growth during early development of white shrimp (*Litopenaeus vannamei*). *Veterinary Microbiology*, 159(3–4), 443–450. <https://doi.org/10.1016/j.vetmic.2012.04.029>
- Pereira, C., Costa, P., Duarte, J., Balcão, V. M., & Almeida, A. (2021). Phage therapy as a potential approach in the biocontrol of pathogenic bacteria associated with shellfish consumption. In *International Journal of Food Microbiology* (Vol. 338). Elsevier B.V. <https://doi.org/10.1016/j.ijfoodmicro.2020.108995>
- Pinoargote, G., Flores, G., Cooper, K., & Ravishankar, S. (2018). Effects on survival and bacterial community composition of the aquaculture water and gastrointestinal tract of shrimp (*Litopenaeus vannamei*) exposed to probiotic treatments after an induced infection of acute hepatopancreatic necrosis disease. *Aquaculture Research*, 49(10), 3270–3288. <https://doi.org/10.1111/are.13791>
- Rico, A., Phu, T. M., Satapornvanit, K., Min, J., Shahabuddin, A. M., Henriksson, P. J. G., Murray, F. J., Little, D. C., Dalsgaard, A., & van den Brink, P. J. (2013). Use of veterinary medicines, feed additives and probiotics in four major internationally

- traded aquaculture species farmed in Asia. *Aquaculture*, 412–413, 231–243. <https://doi.org/10.1016/j.aquaculture.2013.07.028>
- Romano, N., Koh, C. B., & Ng, W. K. (2015). Dietary microencapsulated organic acids blend enhances growth, phosphorus utilization, immune response, hepatopancreatic integrity and resistance against *Vibrio harveyi* in white shrimp, *Litopenaeus vannamei*. *Aquaculture*, 435, 228–236. <https://doi.org/10.1016/j.aquaculture.2014.09.037>
- Tan, C. W., Rukayadi, Y., Hasan, H., Thung, T. Y., Lee, E., Rollon, W. D., Hara, H., Kayali, A. Y., Nishibuchi, M., & Radu, S. (2020). Prevalence and antibiotic resistance patterns of *Vibrio parahaemolyticus* isolated from different types of seafood in Selangor, Malaysia. *Saudi Journal of Biological Sciences*, 27(6), 1602–1608. <https://doi.org/10.1016/j.sjbs.2020.01.002>
- Tsai, C. Y., Santos, H. M., Hu, S. Y., Sang, C. Y., Yanuaria, C. A. S., Lola, E. N. G., Tayo, L. L., Nitura, K. M. A., Liu, C. H., & Chuang, K. P. (2019). LpxD gene knockout elicits protection to *Litopenaeus vannamei*, white shrimp, against *Vibrio parahaemolyticus* infection. *Aquaculture International*, 27(5), 1383–1393. <https://doi.org/10.1007/s10499-019-00398-y>
- Wang, Y., al Farraj, D. A., Vijayaraghavan, P., Hatamleh, A. A., Biji, G. D., & Rady, A. M. (2020). Host associated mixed probiotic bacteria induced digestive enzymes in the gut of tiger shrimp *Penaeus monodon*. *Saudi Journal of Biological Sciences*. <https://doi.org/10.1016/j.sjbs.2020.07.010>
- Wangman, P., Chaivisuthangkura, P., Sritunyalucksana, K., Taengchaiyaphum, S., Senapin, S., Pengsuk, C., Sithigorngul, P., & Longyant, S. (2017). Development of monoclonal antibodies specific to ToxA and ToxB of *Vibrio parahaemolyticus* that cause acute hepatopancreatic necrosis disease (AHPND). *Aquaculture*, 474, 75–81. <https://doi.org/10.1016/j.aquaculture.2017.03.039>
- Xie, J., Bu, L., Jin, S., Wang, X., Zhao, Q., Zhou, S., & Xu, Y. (2020). Outbreak of vibriosis caused by *Vibrio harveyi* and *Vibrio alginolyticus* in farmed seahorse *Hippocampus kuda* in China. *Aquaculture*, 523. <https://doi.org/10.1016/j.aquaculture.2020.735168>
- Zhou, X. xia, Wang, Y. bo, & Li, W. fen. (2009). Effect of probiotic on larvae shrimp (*Penaeus vannamei*) based on water quality, survival rate and digestive enzyme activities. *Aquaculture*, 287(3–4), 349–353. <https://doi.org/10.1016/j.aquaculture.2008.10.046>
- Ziaei-Nejad, S., Rezaei, M. H., Takami, G. A., Lovett, D. L., Mirvaghefi, A. R., & Shakouri, M. (2006). The effect of *Bacillus* spp. bacteria used as probiotics on digestive enzyme activity, survival and growth in the Indian white shrimp *Fenneropenaeus indicus*. *Aquaculture*, 252(2–4), 516–524. <https://doi.org/10.1016/j.aquaculture.2005.07.021>

Chapter 7

INSECT-BASED AQUAFEEDS: A SUSTAINABLE SOLUTION FOR FISH NUTRITION

R. M. Sharven¹ and Jiun-Yan Loh^{1,2}*

¹ Faculty of Applied Sciences, UCSI University, 56000 Cheras, Kuala Lumpur, Malaysia.

² Centre of Research for Advanced Aquaculture (CORAA), UCSI University, 56000, Cheras, Kuala Lumpur, Malaysia

*Corresponding author: lohjy@ucsiuniversity.edu.my

ABSTRACT

In routinely used in aquaculture, insect meals have been proven in many studies to be able to replace fish and soya bean meals partially or totally. However, the amino acid and fatty acid composition, as well as the digestibility of various insect diets is insufficient. Hence, research is needed to evaluate the protein, amino acid, lipid, and fatty acid content of insect meal for comprehensive fish nutrition. In general, insect meal contains more essential amino acids than other meal types, with amounts that are like fishmeals, including methionine, which can be a good supply of the essential nutrient. Insects also rich in lipids and fatty acids, and some antimicrobial components like chitin, which is able to inhibit against pathogenic microorganisms, and further enhance animal health. This chapter discusses the beneficial functions of insect based aquafeed for aquaculture

KEYWORDS: Aquafeeds, insect-based feed, marine organisms, protein, feed replacement

7.1 Introduction

The existing global agricultural production system confronts several obstacles in meeting the rising need of animal byproducts, such as livestock, is expected to increase to 58% by 2050 compared to 2010. A revolution within the variety of animal protein production is highly expected to meet the global demand. As a result of the increased need, food resources, particularly the components used in the formulation of feed, may be put under greater strain in aquaculture feed. Conventional ingredients like soybean,

wheat, corn, and animal meal are projected to increase in terms of price in the upcoming years (Freccia, *et al.*, 2020), this could affect the production cost of aquaculture feed, and the challenges will remain for the next decades. The intensive aquaculture system has been growing in popularity and has emerged as the most promising and practical way to supply the global demand for cultured species. Insect meals offer a novel nutritional source that allows for the highest proportion of fishmeal substitution without compromising the cultivated species' development performance. The fast-growing, immune-enhancing capability and good nutritional quality of insect meals bring probable benefits for cultured species in growth and health promotion (Henry *et al.*, 2015).

Fishmeal is processed through by-catch fishery. According to a statistic, global total landings in 2009 were 22.8 million tonnes of non-food fish (25.7%). Of this volume, 17.9 million tonnes (20.2%) of this total were converted into fishmeal and oil meal (FAO, 2012). Fishmeal and fish oil output in the globe has maintained at around 5.0–6.0 million tonnes per year. The soya bean harvest is by far the most widely grown in the world, increasing by 1.4 % from 2010 to 2011. Soya bean output reached 210.9 million tonnes in 2009, representing for slightly more than half of global oilseed meal production (FAO, 2012). Nogales Mérida *et al.* (2018) found that omnivore and carnivorous fish require higher protein in their diets than the herbivorous or detritivores species. Carnivorous fish require between 55% and 45% crude protein (CP), whereas omnivorous fish require between 45% and 35% CP; these ranges contrast with those of other monogastric vertebrates, which require between 12% and 30% crude protein (CP) (Wilson, 2002). Fish protein needs are directly tied to their amino acid profiles, which are determined by each species' specific amino acid composition.

Fishmeal production output has declined as the marine environment has worsened and fisheries have been depleted, and the price has risen from US\$ 600 per metric tonne in 2005 to US\$ 2000 per metric tonne in June 2010. During the year 2010, the International Monetary Fund (IMF) expects the increased trend in price to continue, with economic implications for animal production. This required for ensuring the need of renewable protein sources, which are especially important in fish diets since fish require a high amount of protein, either from fish feed or perhaps another source (Manzano-Agugliaro *et al.*, 2012).

Due to the ecological challenges (deforestation, stock depletion in fishery products, fluctuating cost) caused by the excessive use/over-dependence of fishmeal in aquafeed (SánchezMuros *et al.*, 2014, Mousavi *et al.*, 2020). Recurrent environmental concerns, as well as fluctuating fishmeal (FM) production and prices, it has prompted the marine feed sector for an alternate source of protein to meet the growing need for aquafeed in the aquaculture sector. Insect-based aquafeeds, often known as insect meal (IM), have garnered significant international interest as a replacement for FM. Important feeding experiments have shown its potential value not just as a protein replacement, but also as a source of bioactive compounds that boosts immunostimulatory actions in a variety of marine species.

Generally, aquafeeds are made using grains, oilseeds, and marine-origin substances, which may act as an antidote to the current emerging issue. Aquafeeds must also compete for feed components with

terrestrial animal industries as pet, chicken, and swine, and complementing ingredients must be developed. Many research have concentrated on alternative sources of protein, but the benefits of plant proteins, microorganism-based proteins, and various animal by-products are currently being addressed some limitations, however significant drawbacks, such as anti - nutritional factors and an uneven nutritional profile, are now being addressed. Throughout this sense, using invertebrates to feed aquatic animals might be another alternative and a critical component of a solution.

The introduction of innovative nutritional sources in commercial aquafeed represents a crucial research field in the sector of aquaculture. Insects have great potential to be nutritious, immune-boost, and sustainable aquafeed that should be tested more in the future. These new sources also offer other advantages as their nutritional value and anti-microbial effect can be modified through intensive culture systems or even with genetic manipulation (Sánchez-Muros *et al.*, 2014).

In this case, using insects as a protein supplement for livestock feed is a realistic choice. Insect production on a commercial scale might provide consistent supply in terms of quantity, quality, and price. The use of insect proteins as feed ingredients in aquaculture was permitted by the European Commission in 2017. The revision to the processed animal protein regulations addresses the horizontal slaughterhouse provision for domestic livestock, which also applies to insect farms and mandates that processed protein be killed in a legally designated slaughterhouse before it can be used in aquafeed. Only 'feed grade' substrates, such as vegetal origin materials or a restricted number of animal origin materials, such as fishmeal, non-ruminant blood products, egg and egg products, milk and milk-based products, honey, and rendered fats, can be given to these insect species (Invertebrate IT, 2021). Some species that are approved by the new EU Regulation 2017/893 entries into force as of 1st July 2017 are as shown in Figure 7.1.

7.2 Benefits of insect meals in terms of physiological improvement and immunity enhancement

Insect meal (IM) has already begun receiving international interest as a potential alternative for FM. Significant feeding studies have demonstrated its potential use not only as a protein substitute but also as a functional element that increases immunostimulatory effects in various aquatic species. In aquatic species, the functional features of IMs, also include their immunomodulatory and physiological effects. Biological metrics such as peroxidation, serum biochemistry, haematological qualities, and immune-related genetics are examples of important factors to examine. In short, low levels of IM supplementation might help specific aquaculture species increase their survival (bivalves, crustaceans etc), immunity, and feed conversion efficiency.

The most important considerations in deciding the choice of IM for aquafeed are nutritional profiles, notably protein and fat levels, nutrient absorption, and insect production. However, the nutritional levels of IM, such as elements (K, Na, Ca, Cu, Fe, Zn, Mn, and P), and vitamin B group vitamins, vitamins A, D, E, K, and C, vary according on sources of food, organisms, and processing techniques. The immunostimulatory effects resulting from the IM are often interpreted in a variety of ways. According to

the research, these health-promoting advantages are largely attributed to either direct generation of antimicrobial peptides (AMPs) or indirect transport of chitin or other bioactive chemicals. Furthermore, key molecules in IM, like as AMPs and chitin, might be employed as functional components, allowing the immune-stimulating and antioxidant properties to be fully investigated in order to improve the general health of the aesthetic species. In addition, the immune-stimulating and antioxidant properties of AMPs in insects might make them a natural option to pharmaceutical medications and antibiotic in aquaculture. To understand the functions of AMPs and their consequences on fish physiological and immunological functioning indices in the attempt of natural alternative exploitation, further genetic-based research on screening and extracting AMP genes in IM are needed. To validate the efficacy of IM in illness suppression and management, field studies and in vivo evaluations are required (Mousavi *et al.*, 2020).



Figure 7.1 House cricket (*Acheta domestica*) (up left); Yellow mealworms(*Tenebrio molitor*) (up right); black soldier larvae (*Hermetia illucens*) (down left); house cricket (*Acheta domestica*) (down right); Common housefly larvae (*Musca domestica*) (Indiviglio, 2022; Indiviglio, 2022; Saanvi, 2021; Heuzé and Tran G, 2015) and other insects like the lesser mealworm (*Alphitobius diaperinus*), banded cricket (*Gryllodes sigillatus*) and the field cricket (*Gryllus assimilis*)

According to certain studies, chitin stimulates the cell-mediated and systemic immune responses of fish, potentially boosting their resistance to infectious illnesses (Loh, 2017). Immune system response peptides present in IMs, such as AMPs, aid in the protection of the host against bacterial infection. Environmental factors, insect species, nutritional balance in meals, and even the sexes of the insects all influence the kind, amount, and quality of produced AMPs.

Insects have also been shown to have considerable pharmacological capabilities in increasing innate immunity in various species, including humans, animals, and poultry, according to studies. It's also important to understand and study that insects have shown assuring results in treating severe human illnesses such as osteomyelitis, etc.

To study the viability and applicability of FM replacement or IM supplementation in aquafeeds, it is critical to first grasp the physiological and immunomodulatory processes that underpin the concept. The amount of ethereal extract in insect meal varies, and this affects the crude energy (kcal kg^{-1}) of the diets, the energy protein ratio, and the aethereal extract of the carcasses. Insect meals with high levels of aethereal extract were not included as frequently in the diets. High levels of aethereal extract are conducive to fat oxidation (rancidity), which shortens the storage life of this product. It is suggested that antioxidant supplements be used in insect feeds. A short-term option would be to improve pre-processing and production techniques to remove the surplus lipid in the meals and use it as a lipid source in feeds or other industries, similar to how domestic animal by-product meals are made (Arru *et al.*, 2019).

Insects may be utilised as a high protein and energy source. Insects also hold a good number of microbial properties. Powdered mealworm has proven to hold microbial properties, while powdered house cricket was characterised by higher microbial diversity since the *Pseudomonas* group was also detected. (Messina *et al.*, 2019). Insect larvae have greater aethereal extract levels because they are storing energy for transformation. Its fatty acid composition is very changeable, indicating that when feed is consumed, the insects' fatty acid profile is modulated, maybe as a precursor to the incorporation of lower-quality EPA (eicosapentaenoic acid, $20:5n^{-3}$) and DHA (docosahexaenoic acid, $22:6n^{-3}$) components. Insect farming helps to natural nutrient recycling and might be a source of high-quality animal protein produced using environmentally friendly technologies.

7.2.1 Chitin as a fibre source

Insect chitin is classified as fibre in general because of its cellulose-like structure. Chitin fibres make up most of an arthropod's exoskeleton (cuticle), and chitin is a polysaccharide made up of glucosamine and N-acetylglucosamine, both of which contain nitrogen atoms (Figure 7.2). Due to the tight coupling of proteins in chitin fibres and variations in chitin according to life stage, estimates of the quantities of chitin and non-digestible protein in insect cuticles vary: hard cuticles have a high protein level of 70–85 percent (dry weight) and a low chitin concentration of 15–30 percent, whereas soft cuticles have a 50 percent chitin and protein content (Chapman 2013; Jonas-Levi & Martinez., 2017).




Figure 7.2 Chemical representation of the cellulose like structure of chitin Merzendorfer (2003).

According to Finke (2007), the digestibility of proteins/AAs in insects depends on how much of the AAs are bound to chitin or scleroprotein, which is typically present in adult insect cuticles (Becker & Yu, 2013). Many fish species with 'natural diets,' such as shrimp, crabs, crayfish, and insects, have chitinolytic action from enzymatic reactions like chitinase in the abdomen and chitobiose in the intestinal wall (Lindsay, 1983). These proteins or amino acids can be available for fish nutrition due to chitinolytic activity from enzymes like chitinase in the abdomen and chitobiose in the internal consistency reliability (Gutowska *et al.*, 2004).

These chitinase activities were responsible for a reduction in *Escherichia coli* and caecal *Salmonella sp.* and an increase in *Lactobacillus acidophilus* in broilers fed chitin from shrimp meal (Khempaka *et al.*, 2011). Sea bass fed varied levels of chitin (0, 0.5, 1, 2, 3, and 4 g kg⁻¹) showed enhanced growth and feed efficiency up to 2 g kg⁻¹ of chitin. Similarly, adding chitin to sea bass fingerlings' meals increased the length of their villus and microvillus in the intestine (Zaki *et al.*, 2015). However, as mentioned by Sánchez-Muros *et al.* (2016), the amount of chitin in the diet must be examined since too much chitin might promote weight gain.

7.3 Insect farming for aquafeed

Use of insect meals in animal nutrition demands large-scale insect rearing to provide a steady supply of insects for animal feed. Insect farming is difficult because insects have specific environmental (temperature and humidity), feeding, and population demands, especially during reproduction (Leppla, 2002). Cultivation of insects can be partial (from egg to larva, nymph to adult), entire (from egg to adult), or full (from egg to adult) (egg-egg). The easiest insects to grow are small, multivoltine herbivores that are terrestrial and have simple environmental requirements, such as pest species of crops, stocked items, or gardens (Leppla, 2002). This method of mass raising has become established, particularly for silk production, fishing bait, and pet food (Schabel, 2010).

Recently, Singaporean-based agri-food tech company (Nutrition Technologies Sdn Bhd) has opened their industrial-scale black soldier fly (BSF) farming factory in Malaysia, to develop insect protein. The company has been initiated by building a large network of factories in the region to supply high-quality IM for the aquaculture, pet food, and animal feed industries, which primarily supports circular economy

practices throughout the supply chain. More and more companies like ENTOFOOD Sdn Bhd have focused on this insect protein production as an opportunity in establishing a biological process that converts a vast quantity of organic waste into insect biomass and fertiliser every day through the work of detritivorous insects. INSEACT, a Singapore-based company is investing efforts by processing palm oil waste, feeds this to insects and processes the insects into premium ingredients for aquaculture. Life Origin, a social enterprise in Malaysia also focuses on generating positive impact to the environment. They produce environmentally friendly and sustainable protein using food waste and disposable media. These growing number of new start-ups show that the scope for the insect protein as an alternative source would be a major solution towards the increasing demand.

7.4 Constraints and future perspectives on aquafeed

Animal meal and soya bean flour, as well as their derivatives and by-products, are some of the extensively utilised protein sources as feed components. Both are linked to ecological (natural resource mistreatment), economic (feed price fluctuations), or production (quality and quantity variations) issues. Besides the value and dependable commercial-scale manufacturing, one of the key problems with IM inclusion in aqua diets is the nutritional profile range.

Excess fat, amino acid imbalances, mycotoxins, and perhaps antinutritional compounds like chitin are all potential limitations. The immune system and survival rates of animals may be compromised by these individual or cumulative factors. These microscopic organisms, on the other hand, are typically grown in industrial facilities and do not require large amounts of area, particularly when compared to other crops like soybeans. Insects use aids in natural nutrient recycling and provides a high-quality animal protein source made with ecologically friendly methods. The fact that insects are easy to grow and multiply, have a high feed conversion efficiency, and may hunt on organic wastes are all factors in favour of their employment (Mérida *et al.*, 2018). Furthermore, research and actual industry demonstrations have proven that supplying big scale insects and using them as a sustainable protein alternative in the diet of marine aqua creatures is technically viable. Once very little animals can be mass produced in large numbers, their costs can be reduced, and they can be used to replace the traditional protein sources in animal feed.

Are insects in aquaculture nutrition a future eco-friendly approach or a commercial reality? It differs according to the industries. As an example, in the salmonid industry, one of the largest and high-value chains in the aquaculture sector, the insect meal already offers an alternative to fishmeal and soya in the early stages of salmon production. The diets were normally made up of the larvae of the black soldier fly (Freccia *et al.*, 2020). These feeds are in use as an EU- approved commodity, and up-to-date surveys show that Norwegian consumers are positive about eating salmon that have had insect meal within the feed formulation. In the future, an academic approach focused on the blue economy will play a significant role in extending consumer acceptance.

7.5 Conclusion

Although insect feed has been examined for use in animal feed, the results have been mixed. Commercial IM production must be produced to match realistic aquaculture market to move an ecologically friendly strategy to the next level. This is also to meet the need for feed meal; enterprises must collaborate with manufacturers that want to enter the commercial market. Furthermore, understanding the nutritional advantages of insects needs research into insect nutrition, since we may be able to change nutrient levels such as EPA and DHA, as well as some essential amino acids, e.g. lysine, methionine, threonine, and tryptophan ratios, based on nutrient requirement of the targeted species. Better understanding is required such as the presence of mycotoxins, lipid rancidification, or unknown antinutritional effects for IM inclusion feed. This field's knowledge is still in its infancy. It is crucial to focus on the separated components from IMs to better understand the process of immunomodulation. The possibilities for using IM in animal nutrition are quite promising, and novel nutritional methods are only being started.

References

- Arru, B., Furesi, R., Gasco, L., Madau, F., & Pulina, P. (2019). The Introduction of Insect Meal into Fish Diet: The First Economic Analysis on European Sea Bass Farming.
- Becker PM, Yu P (2013) What makes protein indigestible from tissue-related, cellular, and molecular aspects? *Molecular Nutrition and Food Research* 57: 1695–1707.
- Chapman RF (2013) Chapter 16 Integument. In: Simpson SJ, Douglas AE (eds) *The Insects, Structure and Function*, 5th edn, pp. 463–500. Cambridge University Press, Cambridge, UK.
- FAO, 26e30 March 2012. Report of the Sixth Session of the Sub-Committee on Aquaculture.
- Finke MD (2007) Estimate of chitin in raw whole insects. *Zoo Biology* 26: 105–115.
- Freccia, A., Sergio Bee Tubin, J., Nishioka Rombenso, A., & Gustavo Coelho Emerenciano, M.(2020). Insects in Aquaculture Nutrition: An Emerging Eco-Friendly Approach or Commercial Reality?. EmergingTechnologies, *Environment And Research For Sustainable Aquaculture*.
- Freccia, A., Sergio Bee Tubin, J., Nishioka Rombenso, A., & Gustavo Coelho Emerenciano,
- Gutowska MA, Drazen JC, Robison BH (2004) Digestive chitinolytic activity in marine fishes of Monterey Bay, California. *Comparative Biochemistry and Physiology Part A: Molecular and Integrative Physiology* 139: 351–358.
- Henry, M., Gasco, L., Piccolo, G. and Fountoulaki, E., 2015. Review on the use of insects in the diet of farmed fish: past and future. *Animal Feed Science and Technology*, 203, pp.1-22.
- Heuzé V., Tran G., 2015. Housefly maggot meal. Feedipedia, a programme by INRAE, CIRAD, AFZ and FAO. <https://www.feedipedia.org/node/671> Last updated on October 21, 2015.
- Indiviglio, F., 2022. Prepared Diets and Food Supplements for House Crickets - Product Review. [online] That Reptile Blog. Available at: <<http://blogs.thatpetplace.com/thatreptileblog/2008/12/05/prepared-diets-and-food-supplements-for-house-crickets-product-review/#.YntcC-hBy3A>> [Accessed 11 May 2022].
- International Monetary Fund, 2010. International Monetary Fund Primary Commodity Prices. <http://www.imf.org/external/data.htm>.
- Invertebrate IT. 2021. New EU regulation authorising the use of insects' proteins in aquaculture - INvertebrateIT. [online] Retrieved from: <http://invertebrateitproject.eu/2017/07/03/new->
- Jonas-Levi A, Martinez JJI (2017) The high level of protein content reported in insects for food and feed is overestimated. *Journal of Food Composition and Analysis* 62: 184– 188.
- Khempaka S, Chitsachapong C, Molee W (2011) Effect of chitin and protein constituents in shrimp head meal on growth performance, nutrient digestibility, intestinal microbial populations, volatile fatty acids, and ammonia production in broilers. *Journal of Applied Poultry Research* 20: 1–11.
- Leppla, N.C., 2002. Rearing of insects. In: Resh, V.H., Carde, R. (Eds.), *Encyclopedia of Insects*. Academic Press, San Diego, California, pp. 975e979.
- Lindsay GJH (1983) Chitinolytic enzymes and bacteria in fish and their role in nutrition (PhD Thesis). Aberdeen University, United States.
- Loh, J. Y. (2017). The Role of Probiotics and Their Mechanisms of Action: An Aquaculture Perspective. *World Aquaculture*.
- M. (2020). Insects in Aquaculture Nutrition: An Emerging Eco-Friendly Approach or Commercial Reality?. EmergingTechnologies, *Environment And Research For Sustainable Aquaculture*.
- Manzano-Agugliaro, F., Sanchez-Muros, M.J., Barroso, F.G., Martínez-Sánchez, A.,Rojo, S., Pérez-Bañón, C., 2012. Insects for biodiesel production. *Renew. Sustain.Energy Rev.* 16, 3744e3753.
- Mériada, S., Gobbi, P., Józefiak, D., Mazurkiewicz, J., Dudek, K., Rawski, M., Józefiak, A. (2018). Insect meals in fish nutrition. *Reviews in Aquaculture*.
- Merzendorfer, H. (2003). Chitin metabolism in insects: structure, function and regulation of chitin synthases and chitinases. *Journal of Experimental Biology*, 206(24), 4393– 4412.

- Messina; Gaglio, ; Morghese, ; Tolone, ; Arena, ; Moschetti, ; Santulli, ; Francesca, ; Settanni, (2019). Microbiological Profile and Bioactive Properties of Insect Powders Used in Food and Feed Formulations. *Foods*, 8(9), 400–. doi:10.3390/foods8090400
- Mousavi, S., Zahedinezhad, S. and Loh, J.Y., 2020. A review on insect meals in aquaculture: The immunomodulatory and physiological effects. *International Aquatic Research*, 12(2), pp.100-115.
- Mousavi, S., Zahedinezhad, S. and Loh, J.Y., 2020. A review on insect meals in aquaculture:
- Nogales-Mérida, S., Gobbi, P., Józefiak, D., Mazurkiewicz, J., Dudek, K., Rawski, M., Kierończyk, B. and Józefiak, A., 2018. Insect meals in fish nutrition. *Reviews in Aquaculture*, 11(4), pp.1080-1103.
- Sánchez-Muros M, De Haro C, Sanz A, Trenzado CE, Vilareces S, Barroso FG (2016) Nutritional evaluation of *Tenebrio molitor* meal as fishmeal substitute for tilapia (*Oreochromis niloticus*) diet. *Aquaculture Nutrition* 22: 943–955.
- Sánchez-Muros, M.J., Rentería, P., Vizcaino, A. and Barroso, F.G., 2020. Innovative protein sources in shrimp (*Litopenaeus vannamei*) feeding. *Reviews in Aquaculture*, 12(1), pp.186-203.
- Schabel, H.G., 2010. Forests insects as food: a global review. In: Durst, P.B., Johnson, D.V., Leslie, R.N., Shono, K. (Eds.), *Forests Insects as food: Humans Bite Back*. FAO, Bangkok, Thailand, pp. 37e64.
- The Business Times. 2021. Nutrition Technologies raises US\$5m, opens insect protein factory in Malaysia. [online] Retrieved from: <https://www-businesstimes-com-sg.cdn.ampproject.org/c/s/www.businesstimes.com.sg/garage/nutrition-technologies-raises-us5m-opens-insect-protein-factory-in-malaysia?amp>
- The immunomodulatory and physiological effects. *International Aquatic Research*, 12(2), pp.100-115.
- Wilson RP (2002) Amino acids and proteins. In: Halver JE, Hardy RW (eds) *Fish Nutrition*, pp. 144–175. Academic Press, San Diego, CA.
- Zaki MA, Salem ME-S, Gaber MM, Nour AM (2015) Effect of chitosan supplemented diet on survival, growth, feed utilization, body composition and histology of sea bass (*Dicentrarchus labrax*). *World Journal of Engineering and Technology* 3: 38–47.



e ISBN 978-967-2782-28-5



